

## Section 24(c) Special Local Need Label

# RESTRICTED USE PESTICIDE (GROUND AND SURFACE WATER CONCERNS)

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

THIS PRODUCT IS A RESTRICTED-USE HERBICIDE DUE TO GROUND AND SURFACE WATER CONCERNS. USERS MUST READ AND FOLLOW ALL PRECAUTIONARY STATEMENTS AND INSTRUCTIONS FOR USE IN ORDER TO MINIMIZE POTENTIAL FOR ATRAZINE TO REACH GROUND AND SURFACE WATER.

### FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF KANSAS

## CORNBELT ATRAZINE 4L HERBICIDE

EPA Reg. No. 11773-1 EPA SLN No. KS-040003

FOR WEED CONTROL IN ROW CROP STUBBLE (SOYBEANS, CORN, OR SORGHUM) FOLLOWING HARVEST ON GROUND TO BE ROTATED TO CORN OR GRAIN / FORAGE SORGHUM THE NEXT YEAR.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

#### **DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, and precautions including statements pertaining to the Worker Protection Standards, on the EPA-registered Cornbelt Atrazine 4L label.

This label must be in the possession of the user at the time of application.

## Cornbelt Atrazine 4L Applied Alone - Row Crop Stubble To Be Planted To Corn or Sorghum (Grain or Forage) the Next Spring

### Soybeans - Corn or Sorghum - Soybeans Rotation on Medium and Fine Textured Soils

Cornbelt Atrazine 4L may be applied to untilled soybean stubble following soybean harvest for control of fall germinating weeds and control of early spring germinating weeds. Broadcast 3.2 to 4.0 pts./A in a minimum of 10 gals. of spray mixture for ground application, with or without crop oil concentrate, depending on soil texture classification and the percent of soil covered with plant residue. On soils not highly erodible, apply 3.2 to 4.0 pts./A. On highly erodible soils where conservation tillage is practiced, leaving at least 30% of the soil covered with plant residues, apply 3.2 to 4.0 pts./A as a broadcast spray. If the soil coverage with plant residue is less than 30%, a maximum of 3.2 pts./A of Cornbelt Atrazine 4L may be applied. In the following spring if supplemental weed control is needed, up to 1.5 lb. a.i./A of atrazine may be applied as Cornbelt Atrazine 4L (1 pint of Cornbelt Atrazine 4L contains 0.5 lb. atrazine active ingredient) or atrazine in registered prepack products. If weeds are present at application which are not on the product label(s), use a tank mixture of Cornbelt Atrazine 4L with an approved foliar applied herbicide.

Apply from soybean harvest up to December 31 but DO NOT apply to frozen ground. DO NOT use where the soil surface conditions favor wind erosion. To minimize the potential of wind erosion, do not fall till soybean stubble after Cornbelt Atrazine 4L application. In no-till operations, maintain desired crop residues on the surface. Plant corn or sorghum the following spring with minimal soil disturbance.

Observe all applicable use limitations and precautions, including setbacks, on the Cornbelt Atrazine 4L label. *The total amount of atrazine must not exceed 2.5 lbs. a.i. / Acre per calendar year.* DO NOT apply Cornbelt Atrazine 4L in the fall if soybeans are to be planted the next spring.

**Note: DO NOT** plant any crops except corn, grain sorghum or forage sorghum the spring following after harvest Cornbelt Atrazine 4L application or illegal residues may result. **DO NOT** graze treated areas within six months after application or illegal residues may result.

Continuous Corn - Corn, Sorghum - Sorghum, or Alternating Corn / Sorghum Rotations on Medium and Fine Textured Soils:

Cornbelt Atrazine 4L may be applied to corn or sorghum stubble following corn or sorghum harvest for control of fall germinating weeds and control of early spring germinating weeds. *The total amount of atrazine to the corn or sorghum crop plus the after crop harvest application must not exceed 2.5 lbs. a.i. / Acre per calendar year.* 

If no atrazine was applied to the corn or sorghum crop prior to harvest, broadcast 3.2 to 4.0 pts./A in a minimum of 10 gals. of spray mixture for ground application, with or without crop oil concentrate, after harvest depending on soil texture classification and the percent of soil covered with plant residue. On soils not highly erodible, apply 3.2 to 4.0 pts./A. On highly erodible soils where conservation tillage is practiced, leaving at least 30% of the soil covered with plant residues, apply 3.2 to 4.0 pts./A as a broadcast spray. On highly erodible soils where plant residue is less than 30%, a maximum of 3.2 pts./A of Cornbelt Atrazine 4L may be applied. In the following spring if supplemental weed control is needed, up to 1.5 lb. a.i./A of atrazine may be applied as Cornbelt Atrazine 4L (1 pint Cornbelt Atrazine 4L contains 0.5 lb. atrazine active ingredient) or atrazine in registered prepack products. If weeds are present at application which are not on the product label(s), remove them with a shallow field cultivation or use an approved foliar applied herbicide tank-mix partner with Cornbelt Atrazine 4L.

If atrazine was applied in the production of the harvested corn or sorghum, the maximum after harvest rate would be 2.5 lbs. a.i./A MINUS the rate applied to the harvested crop, or the rate above, whichever is least. The total of the spring preemergence and postemergence (if used) application(s) and the after harvest application must not exceed 2.5 lbs. a.i./A of atrazine per calendar year.

Apply from corn or sorghum harvest up to December 31 but DO NOT apply to frozen ground. Ground may be tilled before or after application. DO NOT exceed a 2 to 3 inch incorporation depth if tilled after application. In no-till or minimum tillage operations, maintain desired crop residues on the surface. Plant corn or sorghum the following spring with minimal soil disturbance.

Observe all applicable use limitations and precautions, including setbacks, on the Cornbelt Atrazine 4L label.

Note: DO NOT plant any crops except corn, grain sorghum or forage sorghum the spring following after harvest Cornbelt Atrazine 4L application or illegal residues may result.

DO NOT graze treated areas within six months after application or illegal residues may result.

Van Diest Supply Co. P. O. Box 610 Webster City, IA 50595