

		Post-planting by chemigation	1 – 2 gallons per acre for chemigation applications made after planting when either a pre-plant fumigant product was applied or Majestene was applied in- furrow according to the directions below.	Post-emergence; Apply Majestene through chemigation (trickle, drip or overhead) by injecting Majestene into the irrigation event to allow for incorporation of the product into the soil profile at a depth sufficient to protect the root zone several weeks subsequent to application. Additional applications can be made according to the appropriate degree- day modeling for the nematode species being targeted.
MAJESTENE® has a pre-harvest interval (PHI) of 0 days . Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours .				

Nematode management considerations - To obtain maximum results in a nematode management program, it is important to take soil samples prior to application to assess nematode populations and determination of an appropriate management strategy. Utilization of the services of a field person or crop consultant who is knowledgeable in nematode management is advised.

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

This recommendation must be in the possession of the user at the time of pesticide application.

MAJESTENE® is a registered trademark of Marrone Bio Innovations, Inc.
© Copyright Marrone Bio Innovations, Inc., 2014
Marrone Bio Innovations, Inc.
1540 Drew Ave.
Davis, California 95618
Telephone (530) 750-2800
www.marronebioinnovations.com