

TECHNICAL INFORMATION BULLETIN
 FOR THE USE OF MAJESTENE[®] BIONEMATICIDE
 ON LEAFY VEGETABLES

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(Ee) RECOMMENDATIONS ARE RECOGNIZED.

MAJESTENE[®]
EPA Reg. No. 84059-14

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Crop	Target Nematodes	Application Method	Product Use Rate per Application	Application Instructions
Arugula, Celery, Corn Salad, Cress, Dandelion, Dock, Edible-Leaved Chrysanthemum, Endive, Fennel, Head Lettuce, Leaf Lettuce, Parsley, Purslane, Radicchio, Rhubarb, Spinach, Swiss Chard	Suppression of plant parasitic Root knot nematodes (<i>Meloidogyne incognita</i> , <i>M. javanica</i> , <i>M. arenaria</i> , and <i>M. hapla</i>), Needle nematode (<i>Longidorus africanus</i>), Stunt nematode (<i>Merlinius</i> sp.), Spiral nematode (<i>Rotylenchus</i> sp.) and Dagger nematodes (<i>Xiphinema</i> spp.)	In-furrow or banded over an open furrow during planting Soil drench during transplant In-season chemigation	1 - 2 gallons per acre in-furrow or 1-2% v/v drench or 2 gallons per acre	Apply Majestene at a rate of 1-2 gallons per acre in a directed or banded spray over an open furrow. Drench transplants with a 1-2% v/v dilution of Majestene during transplanting with sufficient water to thoroughly soak the root zone. Apply Majestene at a rate of 2 gallons per acre through overhead irrigation
MAJESTENE [®] has a pre-harvest interval (PHI) of 0 days . Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours .				

Nematode management considerations - To obtain maximum results in a nematode management program, it is important to take soil samples prior to application to assess nematode populations and determine an appropriate management strategy. Utilization of the services of a field person or crop consultant who is knowledgeable in nematode management is advised.

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

This recommendation must be in the possession of the user at the time of pesticide application.



MAJESTENE® is a registered trademark of Marrone Bio Innovations, Inc.

© Copyright Marrone Bio Innovations, Inc., 2017

Marrone Bio Innovations, Inc.

1540 Drew Ave.

Davis, California 95618

Telephone: (530) 750-2800

Toll Free: (877) 664-4476

www.marronebioinnovations.com