

Product Bulletin

E. I. du Pont de Nemours and Company Chestnut Run Plaza, 974 Centre Road Wilmington, DE 19805

Fontelis®

EPA Reg. No. 352-834

Special 2(ee) Recommendation†
(For Distribution and Use in the State of Florida)

For Control of Leafspot (*Cercospora spp*) on Watercress

Refer to product label for Fontelis for general use precautions, mixing directions, application directions, and restrictions.

ATTENTION

†This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Fontelis® fungicide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Fontelis according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Fontelis fungicide.

Directions for Use

| Crop | Target Disease | Use Rate per Acre (fl oz) | Remarks |
|---|---------------------------------------|---------------------------|--|
| Watercress | Leafspot (<i>Cercospora spp</i>) | 24 to 30 fl oz | Begin applications prior to disease development and continue on a 7- to 14-day interval. Use higher rate and shorter interval when disease pressure is high. For applications made to watercress, production fields must be drained of water at least 24 hours prior to application and water must not be reapplied to the field for a minimum of 24 hours following the application. |
| Make no more than 2 sequential applications of FONTELIS® before switching to a fungicide with a different mode of action. Minimum time from application to harvest (PHI) is 0 days. Do not exceed 72 fl oz (0.94 lb active ingredient)/acre per year. | | | |

®™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners

R613-007

Issued: 11/26/19

Initial printing