



EPA Reg. No. 71771-14-2935

For Distribution and Use Only Within the States of Georgia, Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Carolina, North Dakota, Ohio, South Carolina, South Dakota, Texas, and Wisconsin

FOR CONTROL OF DOWNY MILDEW, FROGEYE LEAF SPOT, SCLEROTINIA STEM ROT AND WHITE MOLD IN SOYBEAN

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Before using OBRONA™ in accordance with this recommendation, read and follow all applicable directions for use, restrictions, and precautions on the EPA registered label, as found on the package.
- This recommendation should be in the possession of the user at the time of application.
- This recommendation for use of this product is permitted under Section 2(ee) of FIFRA and has not been submitted to or been approved by EPA.

DIRECTIONS FOR USE

CROP	PEST	RATE (Oz/Acre)	DILUTION (Gal/Acre)	TIMING
SOYBEAN	Downy Mildew ( <i>Peronospora manshurica</i> ) Frogeye Leaf Spot ( <i>Cercospora sojina</i> ) Sclerotinia Stem Rot / White Mold ( <i>Sclerotinia sclerotiorum</i> )	1.0	5-200 by ground  2-50 by air	Begin application at first bloom followed by a second application 21 to 28 days after the first application.
Begin foliar applications 5-7 days prior to disease infection in a tank mix or rotation with TARIAN™ or other fungicides for enhanced disease control and yield increase. Can also be applied in a tank mix with the first post-emergent herbicide.				

Use of this product according to this bulletin has not been reviewed or endorsed by the State of Indiana.

WILBUR-ELLIS® logo, FUNGICIDE® logo and TARIAN® are registered trademarks and OBRONA™ is a trademark of Wilbur-Ellis Company LLC.

