



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

Gem™ 500 SC Fungicide

EPA Reg. No. 264-826

For Use on Tree Nuts

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF CALIFORNIA

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. It is a violation of federal law to use this product in a manner inconsistent with its label. This product bulletin must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling.

ALMOND			
Disease Control	Rate fl. oz./Acre	Application Timing	Notes
Alternaria Late Blight (<i>Alternaria alternata</i>) Anthracnose (<i>Colletotrichum acutatum</i>) Jacket Rot (<i>Botrytis cinerea</i>) Scab (<i>Cladosporium carpophilum</i>) Shot Hole (<i>Wilsonomyces carpophilus</i>) Rust (<i>Tranzschelia discolor</i>)	2.9 to 3.8	Begin applications preventively and continue as needed on a 7- to 14-day interval.	Use the higher rates and shorter intervals when disease pressure is severe.
Disease Suppression	Rate Oz/Acre	Application Timing	Notes
Blossom Blight (<i>Monilinia</i> spp.)	1.9-2.9	Begin applications at bud stage. If conditions are favorable for disease development, apply again at full bloom and at petal fall, or on a 14- to 21-day spray schedule.	Use the higher rates and shorter intervals when disease pressure is severe.
<p>Restrictions: Do not apply more than 15.2 fl. oz. of Gem™ per acre per season. Do not apply Gem™ within 60 days of harvest. To limit the potential for development of disease resistance:</p> <ul style="list-style-type: none">Do not make more than two (2) sequential applications of Gem™. Then alternate to at least an equal number of sequential applications of labeled, effective non-QoI fungicides with a different mode of action.Do not apply more than four (4) applications of Gem™ or other QoI fungicides per season.			

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at www.bayercropscienceus.com

As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

Gem is a trademark of Bayer.

04/16/07