



Corteva Agriscience LLC 9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

800-258-3033

Hemi® SC EPA Reg. No. 62719-545

2(ee) Recommendation[†]

For Distribution and Use Only in the State of NC, SC, and VA

Suppression of Thrips in Peanut

ATTENTION

- † This recommendation is made as permitted under FIFRA 2(ee) and has not been submitted to or approved by the EPA.
- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Read and follow all applicable directions for use, precautions, and limitations on the product label attached to the container for this product.
- Except as specified in this product bulletin, use of Hemi® SC is subject to all use precautions and limitations imposed by the label affixed to the container for Hemi SC.

Directions for Use

Apply 2.5 to 3.0 fl oz of Hemi SC per acre to provide early season suppression of thrips in peanuts. For heavier infestations, use a higher rate in the rate range.

Using Hemi SC is not a replacement for insecticide seed treatments or in-furrow insecticide applications at planting. Early season, foliar applications of Hemi SC to suppress thrips in peanuts should only be made to fields that have already received an insecticide seed treatment or in-furrow insecticide application for thrips control.

Use of an adjuvant may increase the level of thrips control.

®™Trademarks of Corteva Agriscience and its affiliated companies

R260-006 Issued: 12/09/2024 **Initial Printing**