

FIFRA Section 2(ee) RECOMMENDATION

DIMETHOATE GROUP 1B INSECTICIDE

Drexel

Dimethoate 4EC

Systemic Insecticide – Miticide

FOR DISTRIBUTION AND USE ONLY IN:

Colorado, Kansas, Illinois, Indiana, Missouri, Nebraska, North Dakota, Oklahoma, South Dakota and Texas

THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.

(EPA REG. NO. 19713-231)

This recommendation for DIMETHOATE 4EC insecticide is valid until 12/31/2028,
or until withdrawn, canceled or suspended.

ACTIVE INGREDIENT:

Dimethoate* 43.5%

OTHER INGREDIENTS: 56.5%

TOTAL: 100.0%

* This product contains 4 pounds of Dimethoate per gallon.

**KEEP OUT OF REACH OF CHILDREN
WARNING / AVISO**Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CROP	PEST	RATE OF APPLICATION
Corn – Field, Pop	Corn Leafhopper (<i>Dalbulus maidis</i>)	1 pint per acre (0.5 lb. a.i./Ac.)
APPLICATION INSTRUCTIONS AND RESTRICTIONS: <ul style="list-style-type: none"> • Do not apply to Corn during the pollen-shed period if bees are present. • Maximum application rate: 16.0 fluid ounces per acre (0.5 lb. a.i./Ac.). • Maximum total rate per year: 16.0 fluid ounces per acre (0.5 lb. a.i./Ac.). • Apply in a minimum of 3 GPA (aerial) and 10 GPA (ground) for optimal coverage and efficacy. • Application by chemigation system is allowed. • The REI is 48 hours. • Interval between last application and harvest is: 14 days forage and 28 days grain. • PROHIBITION: Workers are prohibited from entering the treated area to perform detasseling tasks for 4 days in non-arid areas and for 15 days in outdoor areas where the average annual rainfall is less than 25 inches per year. 		
DIMETHOATE 4EC insecticide may be mixed with adjuvants, fungicides, or herbicides to provide disease or weed control and insect management in one application.		

Remarks and Precautions: Refer to the label for Drexel Dimethoate 4EC for additional use directions, restrictions, rotational crop intervals and precautions. These use directions must be in the possession of the user at the time of pesticide application.

Manufactured By:

Drexel Chemical Company

P.O. Box 13327, Memphis, TN 38113-0327

SINCE 1972