

PRODUCT INFORMATION BULLETIN

Use of this product according to this Bulletin has not been reviewed
or endorsed by the Office of the Indiana Chemist

GROUP	11	FUNGICIDE
-------	----	-----------

GROUP	3	FUNGICIDE
-------	---	-----------



EPA Reg. No. 66330-409

**Supplemental Recommendation for the Use of Fortix® Fungicide
to control Tar Spot on Corn**

**FIFRA Section 2(ee) Recommendation
FOR USE and DISTRIBUTION IN THE STATE OF
INDIANA ONLY**

DIRECTIONS FOR USE

This recommendation is made as permitted under FIFRA section 2(ee) and has not been submitted to or approved by the U.S. Environmental Protection Agency. Observe the warranty statements and conditions specified on the Section 3 label for **FORTIX® FUNGICIDE (EPA Reg. No. 66330-409)**. It is important to read and observe all restrictions, precautions and other information appearing on the Section 3 product label. This recommendation must be in the possession of the user at the time of application.

**FORTIX Fungicide is recommended
for use on Corn to Control Tar Spot Under FIFRA 2(ee)**

Disease	Product Use Rate (fl oz/A)	Application Directions
Tar Spot (<i>Phyllachora maydis</i>)	5 - 6	<ul style="list-style-type: none">• For optimum results begin applications when disease first appears and continue as needed on a 7 to 10-day interval on field and seed corn.• Use the high end of the use rate when disease pressure is high and conditions are favorable for disease development.• Apply no later than growth state R4 (early dough stage).

RESTRICTIONS:

- Do not apply more than 12 fl oz/A (0.13 lb ai/A fluoxastrobin and 0.17 lb ai/A flutriafol) of product per year.
- Do not apply more than 0.36 lb ai/A fluoxastrobin per year if sequentially applying other products containing fluoxastrobin. Do not exceed a maximum of 2 applications per season for all fluoxastrobin containing products.
- Do not make more than two applications per year. Apply no later than growth stage R4 (early dough stage).
- The minimum retreatment interval is 7 days.
- Do not use an adjuvant after the V8 stage and prior to the VT stage of corn. An adjuvant may be used at any other growth stage.
- Restricted Entry Interval (REI) for detasselling is 5 days. The REI for all other activities is 12 hours.
- Pre-harvest Interval: Do not apply product within 30 days of harvest (grain, seed, forage or stover).

Manufactured For:

**Arysta LifeScience North America LLC
c/o UPL NA INC.
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406**

FORTIX is a trademark of UPL Corporation Limited Group Company
©2020 UPL Corporation Limited Group Company

031116V01/2ee/IN