NOT FOR USE IN THE STATE OF TEXAS



EPA Reg. No. 91234-96

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U.S. Environmental Protection Agency.

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF MT and WY

For Control of labeled pests on Potatoes

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This product bulletin must be in the possession of the user at the time of pesticide application.
- Read the "Limitation of Warranty and Liability" in the label booklet for Seville™ before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

For Potatoes (and other Tuberous Root and Corm Vegetables):

Сгор	Pests ¹	Rate Per Acre Per Application	Instructions
Potatoes (and other	5 14 111		Annh mhan amhida O matata madiida fint
Tuberous Root and	Buckthorn aphid	Foliar	Apply when aphids & potato psyllids first
Corm Vegetables) Arracacha	Green peach aphid	application:	appear, before populations build to damaging levels.
Arrowroot	Melon aphid Potato aphid	2.75 – 3.25 oz./A	levels.
Bean yam	Folato aprilu	2.73 - 3.23 02.7A	Two applications may be needed to control
Cassava:			persistent aphid & potato psyllid populations.
bitter and sweet			porocent aprilla di potatto por illa populatione.
Chayote root			
Chinese artichoke			
Chufa			
Dasheen (taro)			
Edible canna			
Ginger	Potato psyllid ²	3.25 oz./A	
Jerusalem artichoke			
Leren			
Potato			
Sweet potato			
Tanier			
True yam			
Turmeric			

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 5 gals./A when applied by air and a minimum of 10 gals./A when applied by ground. To provide optimum coverage and penetration, add a penetrating type spray adjuvant at the manufacturer's specified rate.

Restrictions:

- **Do not** apply more than 3.25 oz./A per application.
- **Do not** exceed a total of 11.0 oz./A per crop per season.

Allow a minimum of 7 days between applications.

Allow a minimum of 14 days (PHI) between the last application and harvest.

¹Aphid pests listed in the individual crop sections are the species that most commonly infest that crop. However, **Seville™** will control all aphid species listed in the Product Information section of the federal label.

²Suppression only.

Use information in accordance with Section 2(ee) of FIFRA.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read and follow all applicable directions for use, restrictions, worker protection standard requirements, and precautions on the EPA registered main label attached to the container of **Seville™** (**EPA Reg. No. 91234-96**).



Manufactured for:
Atticus, LLC
5000 CentreGreen Way, Suite 100
Cary, NC 27513

Seville™ is a trademark of Atticus, LLC.