



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

LUNA[®] SENSATION

Reduced Application Rate for use in almond

EPA Reg. No. 264-1090

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF ARIZONA AND CALIFORNIA

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. It is a violation of federal law to use this product in a manner inconsistent with its label. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

ALMOND		
Disease Control	Product Rate	Use Information
Brown rot blossom blight (<i>Monilinia laxa</i>) (<i>Monilinia fructicola</i>) Shot hole (<i>Wilsonomyces carpophilus</i>) Anthracnose (<i>Colletotrichum acutatum</i>) Alternaria (<i>Alternaria alternata</i>) Scab (<i>Cladosporium carpophilum</i>) Rust (<i>Tranzschelia discolor</i>) Jacket rot Green fruit rot (<i>Botrytis cinerea</i>) Powdery mildew (<i>Sphaerotheca pannosa</i>) (<i>Podosphaera tridactyla</i>)	5.0 to 7.6 fl oz/acre	Begin fungicide applications preventatively. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals. Blossom blight: Applications should be made at the critical timings for disease control. Typically this begins at pink bud stage (about 5% bloom). If conditions are favorable for disease development, apply again at full bloom and at petal fall. Refer to University and/or extension guidelines for best application timings.
Disease Suppression	Product Rate	Use Information
Hull rot (<i>Rhizopus spp.</i>) (<i>Monilinia spp.</i>)	5.0 to 7.6 fl oz/acre	For best results apply early hull split and again 14-days later. Refer to University and/or extension guidelines for best application timings.
Restrictions: Do not apply more than 27.1 fl oz of LUNA [®] SENSATION per acre per season. Do not make more than 4 applications per season. Do not apply LUNA [®] SENSATION within 14 days of harvest. To limit the potential for development of disease resistance to these fungicide classes, do not make more than 2 sequential applications of LUNA [®] SENSATION or any Group 7 or Group 11 containing fungicide before rotating with a fungicide from a different Group.		

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*