

FIFRA §2(ee) RECOMMENDATION
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF CONNECTICUT, HAWAII, ILLINOIS,
INDIANA, KENTUCKY, MAINE, MASSACHUSETTS, MICHIGAN, NEW HAMPSHIRE, NEW JERSEY,
NEW YORK, OHIO, PENNSYLVANIA, RHODE ISLAND, VERMONT, VIRGINIA, WASHINGTON, &
WISCONSIN



EPA REG. NO: 10163-357

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This product bulletin must be in the possession of the user at the time of pesticide application.
- This labeling contains supplemental instructions that may not appear on the product label. All applicable directions, restrictions, and precautions on the product label are to be followed.
- This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the United States Environmental Protection Agency (EPA).

PRODUCT INFORMATION

EcoSwing is a biofungicide for use on growing crops in agricultural, greenhouse, ornamental and turf use sites to control fungal diseases such as powdery mildew (oidiums), sour rot, and *Botrytis cinerea*. Optimum disease control is achieved when the biofungicide is used according to label directions and applied in a regularly scheduled preventative spray program.

The use of EcoSwing is compatible with the principles of Integrated Pest Management (IPM) and programs that attempt to minimize disease resistance to fungicides. Unlike single-site mode of action fungicides which are at risk from disease resistance, EcoSwing has a multi-site mode of action, and may be used to delay or prevent the development of resistance to single site fungicides. Consult with your Federal or State Cooperative Extension Service representatives for guidance on the proper use of EcoSwing in IPM and resistance management programs.

PRE-HARVEST INTERVAL: EcoSwing has a zero-day pre-harvest interval and can be applied up to and including the day of harvest

APPLICATION DIRECTIONS

CROP	PEST	FIELD RATE	COMMENTS:
Raspberries	Orange Rust (<i>Arthuriomyces peckianus</i>) Yellow Rust (<i>Phragmidium rubi-idaei</i>) Spur Blight (<i>Didymella applanata</i>) Post-harvest Fruit Rot (<i>Botrytis spp.</i> , <i>Alternaria spp.</i> , <i>Colletotrichum spp.</i>)	2 pts/Acre	For best preventative results, this product must be applied at least every 7-14 days. Thorough coverage is essential to achieving adequate control. Apply in a regularly scheduled preventative spray program. The use of EcoSwing is compatible with the principles of Integrated Pest Management (IPM) and programs that attempt to minimize disease resistance to fungicides.

EcoSwing® is a registered trademark used under license by Gowan Company, L.L.C.



Produced For:
Gowan Company, LLC
PO Box 5569
Yuma, AZ 85366-5569
25-03-05