

EPA Reg. No. 10163-6414

For use on Grapes for control of Black Rot (Guignardia bidwellii) east of the Rocky Mountains including Alabama, Arkansas, Connecticut, Delaware, Florida, Georgia, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Nebraska, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, West Virginia and Wisconsin.

FIFRA §2(ee) RECOMMENDATION: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the United States Environmental Protection Agency (EPA).

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- Do not apply this product in any way that will contact workers or other persons; either directly or through drift. Only protected handlers may be in the area during application.
- For any requirements specific to your State or Tribe; consult the agency responsible for pesticide regulation.
- Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this
 product.

Crop	Diseases Controlled	Rate of Gavel 75 DF (lb/Acre)	Application Directions (See Also Directions for Use)
Grapes	Black Rot (Guignardia bidwellii)	2.0 to 2.5 (1.33 to 1.67 lb mancozeb and 0.16 to 0.20 lb zoxamide)	Apply when shoots are $\frac{1}{2}$ to 1-1/2 inches long. Repeat when shoots are 3 – 5 inches long, and again when shoots are 8 – 10 inches long, and then continue at 7-10 day intervals. Do not apply within 66 days of harvest. Do not apply more than 6 applications or apply more than 15 lb (10 lb Al mancozeb, 1.25 lb Al zoxamide) per acre per year.



Produced For: Gowan Company P.O. Box 5569 Yuma, AZ 85366-5569