

Product Bulletin

Corteva Agriscience

9330 Zionsville Road

Indianapolis, IN 46268

Matrix[®] SG

EPA Reg. No. 352-768

2(ee) Recommendation[†]

For Distribution and Use Only in the State of California.

PLUS PINDAR[®] GT FOR IMPROVED WEED CONTROL IN THE STATE OF CALIFORNIA IN LABELED TREE NUT CROPS

ATTENTION

[†] This recommendation is made as permitted under FIFRA 2(ee) and has not been submitted to or approved by the EPA.

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Read and follow all applicable directions for use, precautions, and limitations on the product label attached to the containers for both of these products.

To the extent consistent with applicable law, Corteva and/or affiliates will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by Corteva. User assumes all risks associated with uses not specified by the label or 2(ee) instructions.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

DIRECTIONS FOR USE:

MATRIX[®] SG applied at the rate of 4.0 oz product per acre can be tankmixed with PINDAR[®] GT herbicide (1.5 to 3.0 pints per acre), to help improve overall weed control in labeled tree nut crops.

It is important to use a postemergence/contact herbicide such as glyphosate or other registered postemergence herbicides to control any emerged weeds in order to obtain the best residual weed control from MATRIX[®] SG Herbicide.

Before using MATRIX[®] SG and PINDAR[®] GT Herbicides read and follow all applicable directions, restrictions and precautions on the EPA registered label for each product.

RESTRICTIONS:

- The field re-entry interval (REI) for this tank mix is 24 hours.
- Preharvest interval (PHI) for this tank mix is 60 days.
- Do not apply this tank mix through any type of irrigation system.

TMTrademarks of Corteva Agriscience and its affiliated companies

R628-014

Issued: 10/15/2021

Replaces: R1952