

FENAZAQUIN	GROUP	21A	ACARICIDE
FENAZAQUIN	GROUP	39	FUNGICIDE

**FIFRA §2(ee) RECOMMENDATION  
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF CALIFORNIA**

**Additional Pest for Avocado**



**EPA REG. NO. 10163-322**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.
- This recommendation is made as permitted under FIFRA §2(ee) and has not been submitted to or approved by the EPA.

CROP	PEST	USE RATE FL OZ/ACRE	LB A.I./ACRE	COMMENTS
AVOCADO	Persea Mite (Oligonychus perseae)	24 -36	0.32 – 0.48	<b>Ground Application:</b> Apply in at least 100 gallons of water per acre. Use higher labeled rates for heavier mite pressure. <b>Aerial Application:</b> Apply the labeled rate of Magister SC by air in a minimum of 10 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary
	<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• PHI = 7 days</li> <li>• Do not make more than one application per year.</li> <li>• Do not exceed application rate of 36 fl oz of product (0.48 lb a.i.) per acre per year.</li> </ul>			

**RESTRICTIONS ON ALL CROPS:**

- Do not apply by any type of irrigation system.
- Only ground sprayer, airblast and aerial applications are permitted.
- Do not apply during conditions when product will drift to other crops and non-target areas.
- Do not apply within 100 feet of water bodies (including, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries; and commercial fish farm ponds).
- Do not plant rotational crops, other than those on this label, within 30 days of this application. Do not plant root, tuber and bulb vegetables within 120 days of this application.