

## Supplemental Label

**Rely<sup>®</sup> 280**

Herbicide

### For weed control in tropical and subtropical fruits (medium to large fruit, edible peel) in crop group 23B

This supplemental label expires September 30, 2026, and must not be used or distributed after this date.

#### Active Ingredient:

glufosinate-ammonium\* ..... 24.5%\*\*

Other Ingredients: ..... 75.5%

Total: ..... 100.0%

\* CAS Number 77182-82-2

\*\* Equivalent to 2.34 pounds of active ingredient per U.S. gallon.

EPA Reg. No. 7969-448

WARNING/AVISO

#### Directions For Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Rely<sup>®</sup> 280 herbicide** container label, EPA Reg. No. 7969-448, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Rely 280** before applying.
- Use of **Rely 280** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Rely 280**.

For best results, apply **Rely 280** to emerged, young, actively growing weeds. Warm temperatures, high humidity, and bright sunlight improve the performance of **Rely 280**. Uniform, thorough spray coverage of weeds is required to achieve consistent weed control.

Weed control may be reduced when applications are made to weeds under stress such as drought or cool temperatures. Weeds under stress or in dense populations will require application at the highest specified label use rate. Stressed conditions also include prior treatments of other contact or systemic herbicides. Regrowth of weeds may occur due to the weed stage of growth at application, low use rate, or environmental conditions at the time of application.

#### Product Information

**Rely 280** is a water-soluble nonselective herbicide for application as foliar spray for the control of a broad spectrum of emerged broadleaf and grassy weeds. Refer to the **Rely 280** container label for a complete list of weeds controlled.

<b>Rate of Rely® 280 herbicide</b> (fl ozs/A)	<b>Glufosinate Rate Equivalency</b> (lbs ai/A)
48	0.88
56	1.02
64	1.17
82	1.50
246	4.50

### Application Information

Refer to the **Rely 280** container label for additional application instructions and restrictions.

**Rely 280** may be applied to the following tropical and subtropical fruits with medium to large fruit, edible peel, of crop group 23B:

Achachairu, ambarella, araza, babaco, bilimbi, borojo, cajou (fruit), cambuca, carob, cashew apple, ciruela verde, Davidson's plum, feijoa, fig, (Indian) gooseberry, guava (including cattley, para, purple strawberry, strawberry, yellow strawberry), imbe, imbu, jaboticaba, jujube (Indian), kwai muk, mangaba, Marian plum, mombin (including Malayan and purple), monkeyfruit, nance, natal plum, noni, (mountain) papaya, (Japanese) persimmon, pomerac, rambai, rose apple, Sentul, starfruit, Surinam cherry, tamarind, and uvalha.

### Application Rate and Timing

**Rely 280** may be applied in a single application or in sequential applications.

### Postemergence-directed Application

For postemergence control of weeds present in tropical and subtropical fruits, apply **Rely 280** at 48 to 82 fl ozs/A (see chart below, use rate is dependent on target weed growth size and stage) as a broadcast directed spray anytime during the season up to the day of harvest. **Rely 280** may also be applied as a banded or spot treatment to target emerged weeds. See the **Rely 280** container label for application directions of banded and spot treatments.

Avoid contact of **Rely 280** solution, spray, drift, or mist with green bark, stems, foliage, or fruit as injury may occur to trees. **Only trunks with callused, mature brown bark may be sprayed unless protected from spray contact by nonporous wraps, grow tubes, or waxed containers. Contact of Rely 280 with parts of trees other than mature brown bark can result in serious damage.**

**Sequential applications.** Apply **Rely 280** at a minimum of 30 days apart. Regrowth may occur due to the weed stage of growth at application, low use rate, or environmental conditions. Repeat applications of **Rely 280** may be necessary to control plants generating from underground parts or seed.

<b>Weed Size and Stage</b>	<b>Rely 280 Rate</b> (fl ozs/A)
Weeds < 3 inches in height	48 to 82
Weeds < 6 inches in height pre-tiller grasses	56 to 82
Weeds > 6 inches in height and/or grasses that have tillered	64 to 82

### Crop-specific Restrictions

- **DO NOT** apply more than 82 fl ozs/A of **Rely 280** (1.50 lbs ai/A of glufosinate) in a single application.
- **DO NOT** apply more than a maximum cumulative amount of 246 fl ozs/A of **Rely 280** (4.50 lbs ai/A of glufosinate) from sequential applications per year.
- Maximum number of applications per year: 3
- Separate sequential applications by at least 30 days.
- For spot applications, apply as needed for the desired weed control but **DO NOT** exceed the equivalent of 1.50 lbs ai/A (1.88 fl ozs per 1000 square feet) per application or 4.50 lbs ai/A (5.65 fl ozs per 1000 square feet) from applications per year.
- **DO NOT** apply this product aerially in tropical and subtropical fruits.
- **Pre-Harvest Interval (PHI)** – 1 day.

## Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

*Rely is a registered trademark of BASF.*

© 2024 BASF Corporation  
All rights reserved.

007969-00448.20240214.NVA 2024-04-0598-0037  
Based on: NVA 2021-04-594-0188

BASF Corporation  
26 Davis Drive  
Research Triangle Park, NC 27709

  
We create chemistry