

# Product Bulletin



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Curtail\*

EPA Reg. No. 62719-48

### Advanced Crop Stage Applications in Barley and Wheat for Late Emerging Canada thistle and other Broadleaf Weeds

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Read and follow all applicable use directions, precautions and limitations on the label affixed to the product container for Curtail.

#### Directions for Use

Curtail application is normally recommended from the 4 leaf stage (tillering has begun) up to the jointing stage of barley and wheat.

**To control late emerging labeled weeds such as Canada thistle; Curtail may be applied between the jointing and boot stage of barley or wheat. Please consult the complete label for Curtail for detailed use directions.**

To control or suppress labeled weeds, apply Curtail after maximum emergence of the target weeds but before they exceed 3 inches in height or diameter (for rosettes). To obtain season-long control of perennial weeds such as Canada thistle, apply after the majority of the weed's basal leaves have emerged from the soil up to bud stage. Adjust carrier volumes and application equipment to achieve thorough coverage in denser crop canopies.

The risk of crop injury increases as application timing approaches the boot stage. Do not apply unless the risk of injury is acceptable. Do not apply after the boot stage.

\*Trademark of Dow AgroSciences LLC

D06-033-011

Issued: 06-20-02

Initial printing.