



2(ee) Recommendation

For use in the States of Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, Oklahoma, South Carolina, Texas, and Virginia

Fungicide for Control of Listed Diseases Peanuts

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling.

Do not apply this product in any way that will contact workers or other persons; either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe; consult the agency responsible for pesticide regulation. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

Peanut

Target Disease	Product Use Rate per Application fl oz/A	Use Directions
Early leaf Spot (Cercospora arachidicola) Late leaf spot (Cercosporidium personatum)	2.5 – 5.25 (0.037 – 0.078 lb. ai/A)	Begin applications prior to onset of disease when conditions are favorable for disease development, generally around 30 to 40 days after planting. Reapply DOMARK 230 ME using a 14 day interval. DOMARK 230 ME may be used in State Agricultural Extension advisory (disease forecasting) programs which specify application timing based on environmental factors favorable for disease development. Sufficient water volume must be used to ensure thorough coverage for best disease control. Ground application is recommended for best results. Application may be made by ground, air, or chemigation. Apply in 0.1 to 0.25 inches/A of water for chemigation applications. Chemigation application using excessive water could lead to reduced efficacy.

ID# 18-02-01



EPA Registration No. 80289-7-10163
Manufactured by Isagro S.p.A. for:
Isagro USA, Inc.
430 Davis Drive, Suite 240
Morrisville, NC 27560



Distributed by:
Gowan Company LLC
P.O. BOX 5569
Yuma, AZ 85366-5569