

**DuPont Crop
Protection**

**SPECIAL LOCAL NEED
24(C) LABELING**

**DUPONT™ LANNATE® SP
INSECTICIDE
PUMPKINS
STATE OF CALIFORNIA**

RESTRICTED USE PESTICIDE

DUE TO HIGH ACUTE TOXICITY TO HUMANS.

For retail sale and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. Direct supervision for this product requires the certified applicator to review federal and supplemental label instructions with all personnel prior to application, mixing, loading, or repair or cleaning of application equipment.

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF CALIFORNIA

DUPONT™ LANNATE® SP INSECTICIDE

EPA Reg. No. 352 -342

FOR USE ON PUMPKINS IN THE STATE OF CALIFORNIA

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Location:	State of California
Crop/Site/Commodity:	Pumpkins
Target Pest/Problem:	Cotton melon aphid and black bean aphid
Dosage:	Apply one pound of formulation per acre (0.9 lbs ai/acre).
Dilution Rate:	Ground application - Apply in sufficient water to obtain thorough coverage. Aerial application - Apply in 10-20 gallons of water per acre.
Method of Application:	Ground or air
Frequency/Timing of Application:	Apply when insect populations reach threshold levels. Make applications at 5-7 day intervals. A maximum of three applications may be applied per growing season. Do not apply more than 2.7 lbs ai/acre/crop.
Restricted Entry Interval (REI):	48 hours Do not use in "U-Pick", "Pick Your Own" or similar operation.
Preharvest Interval:	3 days
Other Requirements:	None

Do not apply this material through any type of irrigation system.

LANNATE® SP Insecticide should be used as part of an IRM program that features use of alternate mode of action products, as well as biological, cultural and genetic practices aimed at preventing economic pest damage. LANNATE® SP Insecticide is a member of IRAC Group 1A. After two consecutive applications of LANNATE® SP rotate to an insecticide with a different mode of action (different IRAC group number) before making another application of LANNATE® SP. Application of this product should be based on IPM principles and practices including field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds.

IMPORTANT

BEFORE USING LANNATE® SP, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

(Replaces H-64037)

R-783 050608 05-14-08

For product information call 1-888-6-DUPONT

© 2008 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898