



FIFRA 2(ee) Recommendation
EPA Reg. No. 71512-37-74530
FOR DISTRIBUTION AND USE ONLY IN: TEXAS
THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.

This recommendation for Reviton herbicide is valid until December 31, 2026 or until withdrawn, cancelled, or suspended, whichever comes first.

FOR CONTROL OF VOLUNTEER COTTON

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

CROP	PEST	REVITON RATE OF APPLICATION
None	Volunteer Cotton	1 -2 fl oz/A
DIRECTION FOR USE: <ul style="list-style-type: none"> • Apply to volunteer cotton up to 3 true leaves. • Always include an MSO at 1% v/v. • Aerial applications should be applied with a minimum of 3 to 5 gallons per acre. • Ground applications should be applied with a minimum of 10 to 15 gallons per acre. RESTRICTIONS: <ul style="list-style-type: none"> • Do not apply Reviton at a rate greater than 3 fl oz (0.067 lbs ai) per acre. Do not exceed 6 fl oz (0.134 lbs ai) per acre per year. • Follow all plantback intervals listed on the Reviton label. • Do not allow livestock to graze for 21 days after application. <p>A tank-mix of Reviton + MSO adjuvant will effectively control emerged volunteer cotton resulting from harvest the previous season.</p>		

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED REVITON LABEL MUST BE FOLLOWED.

THIS 2(ee) RECOMMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

(RV011623)

HELM Agro US, Inc.
401 E. Jackson St., Suite 1400
Tampa, FL 33602
Phone: 813.621.8846
Fax: 813.621.0763
info@helmagro.com helmagro.com