

OAT Agrio Co., Ltd.

1-3-1, Kanda Ogawa-machi, Chiyoda-ku

Tokyo 101-0052, Japan

**FIFRA 2(ee) Recommendation†**

**FOR DISTRIBUTION AND USE ONLY IN THE STATES OF WASHINGTON**

# **KALIGREEN**

**EPA Registration No.11581-2**

FOR CONTROL OF POST-HARVEST ROTS (MONILINIA FRUIT ROT, PENICILLIUM SPP., BOTRYTIS GRAY MOLD, AND BULL'S EYE ROT) IN POME FRUIT (SUCH AS APPLES, PEARS, AND QUINCE) WHEN APPLIED WITHIN 7-DAYS PRE-HARVEST

<b>CROP</b>	<b>PEST</b>	<b>RATE</b>
POME FRUITS (Such as apples, pear, and quince)	Post-harvest rots (Monilinia fruit rot, Penicillium spp., Botrytis gray mold, and Bull's eye rot)	2 ½ to 3 lbs/A
Apply within 7-days pre-harvest for best control Pre-Harvest Interval for Kaligreen is 1-day		

<b>IMPORTANT NOTICE</b>
<ul style="list-style-type: none"><li>• †This recommendation is made as permitted under FIFRA 2(ee), as amended and has not been submitted to, or approved by the EPA or state lead pesticide agency.</li><li>• This labeling must be in the possession of the user at the time of application.</li><li>• Read the label affixed to the container of Kaligreen before applying. Carefully follow all applicable use directions.</li><li>• Use of Kaligreen according to this product bulletin is subject to all use precautions and limitations stated above or imposed by the label affixed to the container of Kaligreen.</li></ul>

Kaligreen 2ee 20250820