FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF ARIZONA AND CALIFORNIA

VELUM® ONE

EPA Reg. No. 264-1078

For suppression of Alternaria in Tree Nuts (group 14-12)

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

TREE NUTS (Group 14-12)

African Nut-Tree; Almond; Beechnut; Brazil Nut; Brazilian Pine; Bunya; Bur Oak; Butternut; Cajou Nut; Candlenut; Cashew; Chestnut; Chinquapin; Coconut; Coquito Nut; Dika Nut; Ginkgo; Guiana Chestnut; Hazelnut; Heartnut; Hickory Nut; Japanese Horse-Chestnut; Macadamia Nut; Mongongo Nut; Monkey-Pot; Monkey Puzzle Nut; Okari Nut; Pachira Nut; Peach Palm Nut; Pecan; Pequi; Pili Nut; Pine Nut; Pistachio; Sapucaia Nut; Tropical Almond; Walnut, Black; Walnut, English; Yellowhorn; Cultivars, varieties, and/or hybrids of these.

Disease Suppression	Application Rate	Application Instructions
Alternaria (Alternaria alternata)	6.25 to 6.84 fl oz/acre (0.203 – 0.222 lb/acre fluopyram)	Soil Applications - Apply specified dosage by chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment. For optimum results, apply to newly planted trees or those previously trained to drip, trickle or microsprinkler irrigation. Soil must be lightly pre-wetted to break soil surface tension prior to applications.
		Minimum 30-day interval between soil applications.

Restrictions:

- Maximum single application rate: 6.84 fl oz/acre of VELUM ONE (0.222 lb/acre fluopyram)
- Maximum annual application rate: 13.7 fl oz of VELUM ONE per acre (0.445 lb/acre fluopyram) per year.
- Maximum number of applications per year: 2 (at 6.84 fl oz/acre of VELUM ONE)
- Do not apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).
- · Apply using chemigation equipment.
- **DO NOT** apply VELUM ONE within 14 days of harvest.
- To limit the potential for development of disease resistance to this fungicide class, DO NOT make more than 2 sequential applications of VELUM ONE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after VELUM ONE should be a product from a different FRAC group.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions. For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

Bayer CropScience LP 800 N. Lindbergh Blvd. St. Louis, MO 63167