

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.

## 2(ee) Recommendation:

Use on Tomatoes to control Black mold and Early blight



IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

This Product Bulletin Contains New or Additional Directions For Use Which Are Recommended By Sym-Agro, Inc. Under FIFRA Section 2(ee)

FOR DISTRIBUTION AND USE ONLY IN  
Arizona, California, Georgia, Texas, Florida

EPA REG. No. 84876-2

CROP	PEST	RATE OF APPLICATION
TOMATO (Crop Subgroup 8-10A)	Black mold ( <i>Alternaria alternata</i> ) Early blight ( <i>Alternaria solani</i> )	20.4 – 45.7 fl oz/A (0.34 – 0.75 lb ai/A)
<b>DIRECTIONS FOR USE:</b> Begin applications prior to onset of disease development and continue on a 7 to 10 days interval to maintain disease control. Under conditions of severe disease pressure, use the higher rate. Apply the specified rate as a foliar spray in a minimum of 20 gallons or more of water per acre to assure thorough coverage of plants. Increase spray volume as growth increases in order to provide thorough coverage of foliage and fruit for optimum disease control.		
<b>RESTRICTIONS:</b> Do not make more than two sequential applications of ProBlad® Verde Fungicide before alternating or tank mixing with another labeled fungicide with a different mode of action. Do not make more than 5 applications of ProBlad® Verde Fungicide per harvest cycle. <b>PHI: 1 Day</b>		
<b>ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED PROBLAD VERDE® FUNGICIDE (EPA REG. No. 84876-2) LABEL MUST BE FOLLOWED. THIS RECOMMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.</b>		

ProBlad is a trademark of CEV S.A.  
ProBlad Verde\_2ee\_Tomato Expanded Diseases\_2-14-2023