# TECHNICAL INFORMATION BULLETIN

PYRIPROXYFEN GROUP 7C INSECTICIDE



EPA Reg. No. 93930-77

## FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U.S. Environmental Protection Agency.

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF CONNECTICUT, DELAWARE, MAINE, MARYLAND, MASSACHUSETTS, MICHIGAN, NEW HAMPSHIRE, NEW JERSEY, NEW YORK, NORTH CAROLINA, PENNSYLVANIA, VERMONT, VIRGINIA

#### FOR CONTROL OF WOOLLY APPLE APHID IN APPLES

### **DIRECTIONS FOR USE**

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This product bulletin must be in the possession of the user at the time of pesticide application.
- Read the "Limitation of Warranty and Liability" in the label booklet for Positron™ 35 WSP before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

CROP	PEST	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Apple	Woolly Apple Aphid	4 to 5 oz	Apply in 100 to 400 gals/A with ground equipment. Coverage is essentials for good control. Apply in sufficient water volume for tree size. Use of higher water volum will ensure better coverage.
			Woolly Apple Aphid: Apply from delayed dormant to ½ ince green phenological apple growth stage. Under heavy aphid pressure, use the higher rate of Positron™ 35 WSP.

**Positron™ 35 WSP** may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs.

#### **Restrictions:**

- Do not apply within 45 days of harvest.
- Do not make more than 2 Positron™ 35 WSP applications per growing season.
- Do not exceed 10 oz of **Positron™ 35 WSP** per acre per season.
- Do not apply earlier than 14 days after last Positron™ 35 WSP treatment.
- Regardless of formulation, do not apply more than 0.218 lb pyriproxyfen per acre per season.

