## FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

PYRIFLUQUINAZON GROUP 9B INSECTICIDE



EPA Reg. No. 71711-37

For Use On: Whitefly in Potato

**Application Method: Ground Only** 

State(s): Florida

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling and the EPA approved container label must be in the possession of the user at the time of application. Read the label affixed to the container for PQZ® Insecticide before applying. Use of PQZ Insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for PQZ Insecticide.

Potato		
Pest	Rate/Acre	Directions for Use
Whitefly	2.4 to 3.2 fl oz	Apply when pest populations are beginning to build, before reaching economic thresholds. Consult local agricultural
	(0.035 to 0.047	advisor or state extension service for recommendations.
	lb ai)	
		Minimum Spray Volume: 20 gallons of water per acre
		Maximum Dosage Per Acre Per Year: 4.8 fl oz (0.070 lb ai)
		Maximum Applications Per Year: 2
		Minimum Days Between Applications: 14 days
		Preharvest Interval (PHI): 14 days
		Do not feed tops to livestock.

©2019 Nichino America, Inc. PQZ is a trademark and Nichino America logo is a registered trademark of Nichino America, Inc.



Nichino America, Inc. 4550 Linden Hill Road, Suite 501 Wilmington, DE 19808 888-740-7700