

Bayer CropScience LP P.O. Box 12014 2 T.W. Alexander Drive Research Triangle Park, NC 27709 1-866-99BAYER (1-866-992-2937)

## Serenade ASO®

SERENADE ASO is recommended to suppress Citrus Canker on Citrus

EPA Reg. No. 264-1152

# PRODUCT BULLETIN

## FOR DISTRIBUTION IN FL ONLY

**FIFRA Section 2(ee) Recommendation**: This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

## SERENADE ASO is recommended to suppress Citrus Canker on Citrus.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the applicable directions, restrictions and precautionary statements plus all the other information appearing on the EPA-registered product label.

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation. For use only as described on this label. Not for isolation or deformulation. Do not culture.

Serenade ASO may be tank-mixed with other registered pesticides to enhance plant disease control or suppression. This product cannot be mixed with any product with a prohibition against such mixing. When tank-mixing Serenade ASO with other registered pesticides, always read and follow all use directions, restrictions, and precautions of both Serenade ASO and the tank-mix partner(s). Use of the resulting tank mix must be in accordance with the more restrictive label limitations and precautions. Do not exceed label dosage rates.

- 1. Partially fill the spray tank with clean water and begin agitation.
- 2. Add the specified amount of Serenade ASO
- 3. Finish filling the tank to the volume necessary to obtain the proper spray concentration.

It is critical that the spray solution be agitated during mixing and application to assure a uniform suspension. Do not allow spray mixture to stand overnight or for prolonged periods. Maintain a spray solution pH between 4.5 and 8.5.

### **New Use Recommendation**

CITRUS FRUITS – FOLIAR APPLICATION Includes cultivars, varieties and/or hybrids of these commodities.		
Target Diseases	Rate (qt/acre)	Rate when Tank Mixed (qt/acre)
Citrus Canker* - Xanthomonas spp. *NOT FOR USE IN CALIFORNIA	2-4	1-4

For <u>MEDICAL</u> And <u>TRANSPORTATION</u> Emergencies <u>ONLY</u> Call 24 Hours A Day 1-800-334-7577 For PRODUCT USE Information, Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions. For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

151006-R1 Page 1 of 1