



FIFRA 2(ee) Recommendation

EPA Reg. No. 279-3557

FOR DISTRIBUTION AND USE ONLY IN ALABAMA, ARKANSAS, CONNECTICUT, DELAWARE, KENTUCKY, ILLINOIS, INDIANA, LOUISIANA, MAINE, MARYLAND, MICHIGAN, MINNESOTA, MISSISSIPPI, MONTANA, NEW HAMPSHIRE, NEW JERSEY, NEW YORK, NORTH CAROLINA, NORTH DAKOTA, OHIO, PENNSYLVANIA, SOUTH CAROLINA, SOUTH DAKOTA, TENNESSEE, VERMONT, VIRGINIA AND W. VIRGINIA

THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE US EPA.

This recommendation for Topguard® Fungicide is valid until December 31, 2030, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

For Early Season Disease Protection In Spring and Winter Wheat.

Directions For Use

Apply Topguard® fungicide in the fall and/or early spring from tillering to jointing growth stages (Feekes stages 3-7) to provide protection against early season leaf diseases. Topguard fungicide may be tank-mixed with adjuvants, fertilizer, insecticides (such as **Mustang® MAXX** insecticide) or herbicides (such as **Harmony® Extra SG** herbicide with TotalSol® soluble granules or **Finesse® Cereal** and **Fallow** herbicide) to implement disease, insect and weed management. An application of Topguard® EQ fungicide may be applied at flag leaf (Feekes 8-9) for control of foliar diseases as specified by product labels. Apply in a minimum of 2 gallons per acre for aerial application and 10 gallons per acre by ground application.

CROP	PESTS	RATE OF APPLICATION
Wheat (Spring and Winter)	Common Rust and Stripe Rust (<i>Puccinia spp.</i>) Powdery Mildew (<i>Blumeria graminis tritici</i>) Leaf and Glume Blotch (<i>Septoria spp.</i>) Tan Spot (<i>P. tritici-repentis</i>)	5 - 7 fl. oz./A

RESTRICTIONS:

- Do not apply more than 2 applications per year.
- Do not apply more than 0.228 lb. ai of flutriafol/A per year.
- Restricted entry interval (REI) is 12 hours.
- Preharvest Intervals (PHI): Do not apply within 30 days of harvest when grown for grain, 15 days of harvest when grown for hay, zero (0) day PHI when grown for forage.

R-4968 06132025



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000