

Supplemental Labeling

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268 USA

Fontelis®

EPA Reg. No. 352-834

This supplemental label expires on 04/10/2026, and must not be used or distributed after this date.

For Control of Diseases in Greenhouse Lettuce

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Fontelis® fungicide before applying. Follow all applicable directions, restrictions, PPE requirements and precautions on the label affixed to the container for Fontelis.
- Use of Fontelis according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Fontelis.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Directions for Use

Fontelis is a broad-spectrum fungicide, recommended for control of foliar and soil-borne plant diseases in many crops, and has preventive, curative, and locally systemic activity. Fontelis must be applied in a regularly scheduled protective spray program in rotation with other fungicides. Refer to the product label for Fontelis for a complete list of targeted diseases. Follow all mixing and application instructions in the Product Information section of the product label for Fontelis.

Rate and Application Timing

Crop/Crop Group	Target Diseases	Use Rate per Acre (fl oz)	Remarks
Lettuce, greenhouse	Alternaria leaf spot* (<i>Alternaria sonchi</i>) Cercospora leaf spot (<i>Cercospora</i> spp.) Early blight (<i>Cercospora apii</i>) Gray mold* (<i>Botrytis cinerea</i>) Late blight (<i>Septoria apicola</i>) Powdery mildew (<i>Erysiphe cichoracearum</i>) Rust* (<i>Puccinia</i> spp.) Septoria leaf spot* (<i>Septoria</i> spp.)	14-24 fl oz	Begin applications prior to disease development and continue on a 7- to 14-day interval. Use higher rate and shorter interval when disease pressure is high.
	Lettuce drop (<i>Sclerotinia minor</i> , <i>Sclerotinia sclerotiorum</i>)	16-24 fl oz	
Make no more than 2 sequential applications of FONTELIS® fungicide before switching to a fungicide with a different mode of action. Minimum time from application to harvest (PHI) is 3 days. Do not make more than 3 applications per year at the high rate. Do not exceed 72 fl oz (0.94 lb active ingredient)/acre per year. In New York, do not exceed 61 fl ounces (0.80 lbs ai) per acre per year.			

*Not for use in California

Specific Use Restrictions:

- Use this product only in commercial and farm plantings.
- Do not use for home plantings.
- Do not formulate this product into other end-use products.

™®Trademarks of Corteva Agriscience and its affiliated companies

R613-029
 EPA Accepted: 04/06/23
 Initial Printing