



SUPPLEMENTAL LABELING

**RESTRICTED USE PESTICIDE
DUE TO AVIAN AND AQUATIC TOXICITY**

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

DIAZINON AG500

Insecticide

EPA REG NO. 66222-09

**FOR USE ON FIGS IN CALIFORNIA, WATERCRESS IN HAWAII, AND
FILBERTS IN OREGON AND WASHINGTON**

ACTIVE INGREDIENT: Diazinon:	% By Wt.
O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate.....	48.0%
INERT INGREDIENTS*	52.0%
TOTAL	100.0%

Contains 4 lbs. Diazinon per gallon

*Contains xylene range aromatic solvent

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

DIRECTIONS FOR USE

It is violation of federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of pesticide application. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

Crop	Pest	Rate	Comments
FIGS (California Only)	Vinegar flies- <i>drosophila</i> spp., Dried fruit beetle	Apply 1 pt. per 100 gals. of water not to exceed 200 gals. of water per application	Apply as infestations occur. Repeat application after 7 days if necessary.
	<ul style="list-style-type: none"> Do not apply more than 2 pints of product per acre per application and no more than 6 pints of product per acre per year. To avoid illegal residues, allow a minimum of 5 days between the last application and harvest. 		
FILBERTS (HAZELNUTS) (Oregon and Washington Only)	Filbert Leafroller, Aphids	Apply 1-2 quarts in 250-400 gals. of water per acre	Apply as a thorough cover spray. Repeat application after 14 days if necessary.
	<ul style="list-style-type: none"> Do not apply more than 2 quarts of product per acre per application and no more than 6 quarts per year. To avoid illegal residues, allow a minimum of 21 days between the last application and harvest. 		

Crop	Pest	Rate	Comments
WATERCRESS (Hawaii Only)	Cyclamen Mites	Apply 1 pint per 100 gals. of water per acre.	Apply as a thorough cover spray. Spray should be directed into the crowns and growing tips of plants. If infestation is heavy, make a second application within 10 days after the first application.
	<ul style="list-style-type: none"> • Do not apply more than 1 pint of product per acre per application and no more than 2 pints of product acre per season. • To avoid illegal residues; allow a minimum of 5 days between the last application and harvest 		

Diazinon AG500 (66222-9)(supplemental on figs, filberts, watercress EPA app 5-9-02)(CA app 11-23-04)