

# Product Bulletin

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Transform<sup>®</sup> WG

EPA Reg. No. 62719-625

### Special 2(ee) Recommendation<sup>†</sup>

(Not for use in the states of California, New York and Texas)

### For Control of Leafhoppers in Cherry

Refer to product label for Transform<sup>®</sup> WG for general use precautions, mixing directions, application directions, and restrictions.

#### ATTENTION

<sup>†</sup>This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Transform WG insecticide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Transform WG according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Transform WG insecticide.

#### Directions for Use

| Pests       | Transform WG<br>(oz/acre)        |
|-------------|----------------------------------|
| Leafhoppers | 1.5 - 2.75<br>(0.086 lb ai/acre) |

Apply 1.5 to 2.75 oz of Transform WG per acre to provide control of leafhoppers in cherry. Use a higher rate in the rate range under heavy pest pressure and/or when targeting adult leafhoppers. Treat in accordance with local economic thresholds. Consult your Corteva Agriscience representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

#### Restrictions:

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- **Minimum Treatment Interval:** Do not make applications less than 7 days apart.
- Do not make more than four applications per crop.
- Do not make more than two consecutive applications per crop.
- Do not apply more than a total of 8.5 oz of Transform WG (0.266 lb ai of sulfoxaflor) per acre per year.
- Do not apply this product at any time between 3 days prior to bloom and until after petal fall.

<sup>TM</sup>®Trademarks of Corteva Agriscience and its affiliated companies

R396-277

Issued: 08/19/22

Replaces Labels: R396-273 and R396-274