

FIFRA 24(c) Special Local Need Label (SLN)

For Distribution and Use Only in the State of Hawaii.

For Control of Thrips and Planthoppers in Field Corn Grown for
Seed Production, Research, and Breeding Purposes Only.



TOLFENPYRAD	GROUP	21A	INSECTICIDE
	GROUP	39	FUNGICIDE

EPA Reg. No. 71711-31

SLN No. HI-240001

This labeling expires and must not be distributed or used in accordance with this SLN registration after January 28, 2029.

ENVIRONMENTAL HAZARDS

This pesticide is very highly toxic to fish and aquatic invertebrates. **For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean highwater mark. Do not apply when weather conditions favor drift from treated areas.** Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when cleaning equipment or disposing of equipment washwaters. This product is highly toxic to bees and other pollinating insects exposed to direct treatment or residues on blooming crops or weeds. APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE ON THE TORAC® INSECTICIDE LABEL TO PROTECT POLLINATORS. **Do not apply this product or allow it to drift to blooming crops or weeds if bees or other pollinating insects are visiting the treatment area.** Application must be made at least 8 hours prior to bees foraging.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application. Read the label affixed to the container for **TORAC** Insecticide before applying. Use of **TORAC** Insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **TORAC** Insecticide. New use directions appear on this supplemental labeling that may be different from those that appear on the container label.

APPLICATION RATE CHART FOR TORAC INSECTICIDE

Field Corn Grown for Seed Production, Research, and Breeding Purposes Only		
Pest	Rate/Acre	Use Directions
Planthoppers	17.0 to 21.0 fl oz/acre (0.17 to 0.21 lb ai/acre)	<ul style="list-style-type: none">• Apply by ground only: use a minimum of 40 gallons of water per acre.• Allow at least 14 days between applications.• Preharvest Interval (PHI): 1 day
Thrips	24.0 fl oz/acre (0.24 lb ai/acre)	USE RESTRICTIONS <ul style="list-style-type: none">• Do not apply more than 48.0 fluid ounces (0.48 lb ai) per acre per crop cycle.• Do not make more than 2 applications per crop cycle.• Do not make more than 4 applications per year.• No part(s) of the treated crop shall be used as food or feed.• Do not graze livestock on treated fields.
IMPORTANT: THIS LABEL IS ONLY FOR USE BY AUTHORIZED NICHINO AMERICA PERSONNEL, MEMBERS, OR THEIR GROWERS UNDER THE HAWAII CROP IMPROVEMENT ASSOCIATION AND MAY NOT BE COPIED OR RETRANSMITTED IN ANY FORM. NO PART(S) OF THE CROP TREATED WITH TORAC INSECTICIDE SHALL BE DIVERTED AS FOOD FOR HUMAN CONSUMPTION OR FEED FOR ANIMAL CONSUMPTION.		

©2023 Nichino America, Inc. Torac is a registered trademark of Nichino America, Inc.



Nichino America, Inc.
4550 Linden Hill Road, Suite 501
Wilmington, DE 19808
888-740-7700