

RESTRICTED USE PESTICIDE

Due to Acute Oral, Dermal and Inhalation Toxicity and Risks to Wildlife.  
For retail sale to, and use only by Certified Applicators or persons under the direct supervision of a Certified Applicator, and only for those uses covered by the Certified Applicator's certification.



**FIFRA 2(ee) RECOMMENDATION**

For Control of Cotton Jassid, the Two Spotted  
Cotton Leafhopper, in Cotton



WATER MISCIBLE INSECTICIDE

EPA REG. NO. 5481-448

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF Alabama,  
Arkansas, Arizona, Florida, Georgia, Kansas, Louisiana, Mississippi, Missouri,  
New Mexico, North Carolina, Oklahoma, Tennessee, and Virginia.**

**Directions for Use**

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

This recommendation is made as permitted under FIFRA § 2(ee) and has not been submitted to or approved by the Environmental Protection Agency (EPA).

**THIS BULLETIN MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION.**

Before applying **Bidrin 8** please read the complete label affixed to the product container. All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Use of **Bidrin 8** according to this recommendation is subject to the use precautions and limitations imposed by the label on the container.

**IMPORTANT:** Read the entire Directions for Use and the Limited Warranty and Disclaimer on the EPA-registered **Bidrin 8** label before using this product.

**Use Information**

**APPLICATION DIRECTIONS**

BIDRIN 8 must be used in a closed delivery system (See Engineering Controls).

BIDRIN 8 is a water-miscible product which contains 8 lbs of active ingredient per gallon. For foliage applications, mix the required amount of BIDRIN 8 in sufficient water to provide uniform coverage.

**CHEMIGATION**

Do not apply this product through any type of irrigation system.

USE AND DIRECTIONS FOR APPLICATION ON COTTON LATE SEASON USE (First bloom to 30 days prior to harvest)		
Insect	Dosage (fluid ounces/acre)	Directions for Application
Two Spotted Cotton Leafhopper ( <i>Amrasca biguttula biguttula</i> )	6.0 to 8.0	<b>Do not apply more than 16 fl. oz./ac of BIDRIN 8 (1.0 lb. a.i./ac) during this growth period.</b> Apply by air or ground to give uniform coverage. The minimum application interval between applications is 14 days. Do not graze livestock on treated fields or feed treated gin trash. Do not apply within 30 days of harvest.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the restrictions, precautions and directions for use on all included product labels. Users must follow the most restrictive directions and precautionary statements of each product in the tank mixture. **Bidrin 8** is a restricted use pesticide.

©2025 AMVAC Chemical Corporation is a wholly owned subsidiary of American Vanguard Corporation. All Rights Reserved. AMVAC, Bidrin, and their respective logos are trademarks owned by AMVAC Chemical Corporation.

Registrant:  
AMVAC Chemical Corporation  
4695 MacArthur Court, Suite 1200  
Newport Beach, CA 92660 U.S.A.  
1-888-462-6822