



**Bayer CropScience**

Bayer CropScience LP  
P.O. Box 12014  
2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
1-866-99BAYER (1-866-992-2937)

## VELUM® PRIME

For application at a reduced rate in Stone Fruit

EPA Reg. No. 264-1078

### ***FIFRA 2(ee) Recommendation***

#### **FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF MICHIGAN, OREGON, AND WASHINGTON**

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

#### **DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its label.

#### **STONE FRUITS (Group 12-12)**

**Cherry subgroup:** Capulin; Black Cherry; Nanking Cherry; Sweet Cherry; Tart Cherry; cultivars, varieties, and/or hybrids of these.

**Peach subgroup:** Peach; Nectarine; cultivars, varieties, and/or hybrids of these.

**Plum subgroup:** Apricot; Japanese Apricot; Chinese Jujube; Plum; American Plum; Beach Plum; Canada Plum; Cherry Plum; Chickasaw Plum; Damson Plum; Japanese Plum; Klamath Plum; Plumcot; Prune Plum; Sloe; cultivars, varieties, and/or hybrids of these.

| <b>Pest Suppression</b> | <b>Application Rate</b>  | <b>Application Instructions</b>  |
|-------------------------|--------------------------|--|
| Nematodes               | 6.5 - 6.84<br>fl oz/acre | <p><b>Soil Applications</b> - Apply specified dosage by chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment. For optimum results, apply to newly planted trees or those previously trained to drip, trickle or micro-sprinkler irrigation. Soil must be lightly pre-wetted to break soil surface tension prior to applications.</p> <p>Minimum 30-day interval between soil applications.</p> |

#### **Restrictions:**

- Do not apply more than 13.7 fl oz of VELUM PRIME (0.446 lbs Fluopyram) per acre per year, regardless of formulation or method of application (soil or foliar).
- Apply using chemigation equipment.
- Can be applied the day of harvest.
- For soil application, to limit the potential for development of disease resistance to this chemical class, the first foliar fungicide application after VELUM PRIME should be a product from a different FRAC group.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.  
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

*As with any crop-protection product, always read and follow label directions.  
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*