



**DIREX® 4L HERBICIDE**  
**EPA REG. NO. 66222-54**  
**FOR CONTROL OF MANY ANNUAL AND PERENNIAL**  
**GRASSES AND HERBACEOUS WEEDS**

**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR **DIREX® 4L HERBICIDE** BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for DIREX 4L HERBICIDE and this supplement.

---

**FOR USE ON PEPPERMINT IN CALIFORNIA, IDAHO, OREGON, AND WASHINGTON**

---

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**DIRECTIONS FOR USE**

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This supplemental labeling must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.
- These directions can be found on the currently registered EPA Stamped Label.

**PEPPERMINT**

**California, Idaho, Oregon, Washington:** Apply 0.6 to 0.8 quart per acre on soils having 1 to 2% organic matter. Apply 0.8 to 1.6 quarts per acre on soils having 2.1 to 3.0% organic matter. Apply 1.6 to 2.4 quarts per acre on soils having more than 3.0% organic matter.

Do not apply to stands of mint suffering from stress due to low fertility, drought, winter injury, insects, disease or damage from other herbicides or other causes.

Do not apply to snow-covered or frozen ground, as injury to the crop or poor weed control may result.

Do not apply to sand, loamy soil, gravelly soils, or exposed subsoils. Do not apply to soils that have a high salt content and/or high water table or poor drainage that retards mint root development resulting in a shallow root system. Do not apply to soils having less than 1% organic matter.

**Application Timing:** Apply Direx 4L to established (at least one year) stands of mint during the late winter dormant period or after flaming in the spring prior to the emergence of new growth. Do not cultivate after application.

If weeds are present at time of application, the use of a surfactant at 0.25% v/v or crop oil concentrate at 1.0% v/v may be used to increase the performance of Direx 4L postemergence to weeds.

**Tank Mixes and Sequential Treatments:** Direx 4L can be applied either alone or in a program involving tank mixes and/or sequential treatments with other herbicides and adjuvants providing Direx 4L is not applied to actively growing mint plants.

When using a tank mix with other herbicides, use the lower end of the Direx 4L use rate range unless prior experience indicates it is safe to use higher rates. Tank mixes and sequential treatments with other herbicides can increase the risk of crop injury. When using a certain tank mix or sequential treatment for the first time, limit use to a small area to determine safety before treating large areas.

Read the “**LIMITATION OF WARRANTY AND LIABILITY**” in the label booklet for DIREX 4L HERBICIDE before using.

032213

