

Product Bulletin

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Ridgeback™

EPA Reg. No. 62719-749

Special 2(ee) Recommendation†

(For Distribution and Use in the States of Delaware, Maryland, and Pennsylvania)

For the Control of Mexican Bean Beetle on Soybean

Refer to product label for Ridgeback for general use precautions, mixing directions, application directions, and restrictions.

ATTENTION

†This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Ridgeback insecticide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Ridgeback according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Ridgeback insecticide.

Directions for Use

Apply Ridgeback to Soybean at the rate of 8.6 – 13.8 fl oz/acre to control Mexican Bean Beetle.

Restrictions:

- **Preharvest Interval:** Do not apply within 18 days of harvest.
- **Minimum Treatment Interval:** Do not make applications less than 30 days apart.
- Do not make more than four applications per year.
- Do not make more than two consecutive applications per crop.
- Do not apply more than a total of 41.3 fl oz of Ridgeback (0.3 lb ai of bifenthrin and 0.1 lb ai sulfoxaflor) per acre per year.
- No more than two applications may be made to soybean forage.

™® Trademarks of Corteva Agriscience and its affiliated companies

R467-001

Issued: 02/19/24

Initial