

Supplemental Labeling



Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Fontelis[®] FUNGICIDE

EPA Reg. No. 352-834

This supplemental label expires on April 30, 2027, and must not be used or distributed after this date.

**FOR USE IN CEREAL GRAINS EXCEPT RICE, INCLUDING: BARLEY,
MILLETS, SORGHUMS, AND WHEAT
(Not Registered for Use in California)**

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in possession of the user at the time of application.
- Read the label affixed to the container for Fontelis[®] before applying.
- Use of Fontelis according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Fontelis.

Product Information

Fontelis is a broad-spectrum fungicide, recommended for control of foliar and soil-borne plant diseases in many listed crops, and has preventive, curative, and locally systemic activity. Fontelis must be applied in a regularly scheduled protective spray program in rotation with other fungicides. See directions below for specific crop/disease recommendations.

Fontelis can be applied with commonly used ground equipment, hose-end, pressurized, and greenhouse sprayers, air or chemigation equipment, except as otherwise directed, using sufficient water to obtain thorough coverage of plants. Thorough coverage of all foliage is essential for effective disease control. Maintain agitation during mixing and application to assure uniform product suspension.

Use Restrictions

- Use this product only in commercial and farm plantings.
- Do not use for home plantings.
- Do not formulate this product into other end-use products.
- Do not sell, sell into, distribute, or use in Nassau or Suffolk Counties, New York.
- In New York, do not exceed 61 fl oz (0.80 lb ai) per acre per year.
- Not approved for use in the State of California.

Fontelis fungicide labeled Crop and Crop Groups, Pre-Harvest Intervals, Maximum Single

Application Rates, and Total Rates allowed per year

Crop, Crop group or subgroup with examples [†]	Minimum Time from Application to Harvest (PHI days)	Maximum Rate per Acre per Application fl oz product (lb active ingredient)	Maximum Product per Acre per Year fl oz product (lb active ingredient)
Cereal grains* Barley, Wheat, oats, rye	0 day forage and hay, Do not apply after flowering	24 fl oz (0.31 lb)	48 fl oz (0.63 lb)
Cereal grains* Sorghum	0 day forage 30 days grain and stover	24 fl oz (0.31 lb)	48 fl oz (0.63 lb)

*Not for use in California

[†]Refer to the crop specific direction for use tables to determine exactly which crop species in each crop area or crop group are included.

Use Rates and Application Instructions

Cereal grains* except rice, including: barley, millets, sorghums, wheat, corn

Use directions for specific cereal grains are provided below. Cereal grains may be used for grazing, forage, and/or hay within 0 days after the last application.

Crop/Crop Group	Target Diseases	Use Rate per Acre (fl oz)	Application Instructions
Barley*	Scald (<i>Rhynchosporium secalis</i>) Spot blotch (<i>Cochliobolus sativus</i>)	14 - 24 fl oz (0.18 – 0.31 lb ai/A)	Begin applications prior to disease development and continue on a 7- to 14-day interval, depending on the targeted disease. Use higher rate and shorter interval when disease pressure is high.
RESTRICTIONS: Make no more than 2 sequential applications of Fontelis before switching to a fungicide with a different mode of action. Do not apply after flowering (Feekes 10.5.1). Do not exceed 48 fl oz/acre (0.63 lb ai) per year.			
Millet, pearl* millet, proso* Sorghum (milo)* sorghum spp. (sudangrass and hybrids)*	Rust, common (<i>Puccinia sorghi</i>)	10 - 24 fl oz (0.13 – 0.31 lb ai/A)	Begin applications prior to disease development and continue on a 7- to 14-day interval. Use higher rate and shorter interval when disease pressure is high.
RESTRICTIONS: Make no more than 2 sequential applications of Fontelis before switching to a fungicide with a different mode of action. Minimum time (PHI) between application and grain and stover harvest is 30 days. Do not exceed 48 fl oz (0.63 lb ai)/acre per year.			

*Not for use in California

Crop/Crop Group	Target Diseases	Use Rate per Acre (fl oz)	Application Instructions
Wheat* rye* oats* buckwheat* teosinte* triticale*	Leaf and glume blotch (<i>Stagonospora</i> spp., <i>Septoria</i> spp.) Rust, brown leaf (<i>Puccinia recondita</i> f. <i>sp. tritici</i>) Rust, black stem (<i>Puccinia graminis</i> f. <i>sp. tritici</i>) Rust, stripe (<i>Puccinia striiformis</i>)	10 - 24 fl oz (0.13 – 0.31 lb ai/A)	Begin applications prior to disease development and continue on a 7- to 14-day interval, depending on the targeted disease. Use higher rate and shorter interval when disease pressure is high. To optimize yields in cereals, it is important to protect the flag leaf from foliar diseases. For optimizing yield and flag leaf disease control, apply Fontelis fungicide at Feekes 9, 'flag leaf out.'
	Tan spot (<i>Pyrenophora tritici-repentis</i>)	16 - 24 fl oz (0.21 – 0.31 lb ai/A)	
	Disease suppression Powdery mildew (<i>Erysiphe graminis</i> f. <i>sp.</i> <i>Tritici</i>) Scab (<i>Fusarium</i> spp.)	10 – 24 fl oz (0.13 – 0.31 lb ai/A)	
RESTRICTIONS: Make no more than 2 sequential applications of Fontelis fungicide before switching to a fungicide with a different mode of action. Do not apply after flowering (Feekes 10.5.1). Do not exceed 48 fl oz (0.63 lb ai)/acre per year.			

*Not for use in California

™® Trademarks of Corteva Agriscience and its affiliates

R613-015
EPA accepted 05/06/24
Initial printing