

FORMETANATE
HYDROCHLORIDE

GROUP

1A

INSECTICIDE

RESTRICTED USE PESTICIDE

Due to acute oral toxicity

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

REGISTRATION FOR SPECIAL LOCAL NEED FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF TEXAS

Carzol[®] SP

INSECTICIDE

IN WATER SOLUBLE PACKAGING
For Agricultural Use Only

EPA REG. NO. 10163-265, EPA SLN NO. TX-250005

Expiration Date: 12/31/2030

For the control of thrips in oranges, lemons, tangerines and grapefruit

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions, including statements pertaining to the Worker Protection Standards, on the EPA-registered label for Carzol SP Miticide/Insecticide in Water Soluble Packaging are to be followed.
- This labelling must be in the possession of the user at the time of pesticide application.

Corp	Pest	Application
ORANGES, LEMONS, TANGERINES AND GRAPEFRUIT	Thrips	Apply 0.5 to 1.25 lbs. of CARZOL SP per acre as outside coverage spray when thrips appear.
	Restrictions: <ul style="list-style-type: none">• Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 9 days.• Do not apply more than 1.25 lbs. CARZOL SP per acre in one crop season.• Do not apply after fruit reach a diameter of one inch.• If unharvested grapefruit and Valencia oranges are present from the previous crop, an application may be made to the new crop. However, a preharvest interval of 30 days must be observed for the unharvested crop.• Note: If soft brown scale is present in the grove, a suitable scallicide should be applied after the CARZOL SP application.• Aerial applications to orchard crops is prohibited.	

Gowan[®]

24(c) Registrant:
Gowan Company, LLC
P.O. Box 5569
Yuma, AZ 85366-5569