



# FIFRA 2(ee) Recommendation

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF  
ARIZONA AND CALIFORNIA**

## Velum<sup>®</sup> One

**EPA Reg. No. 264-1078**

**For Suppression of Red Leaf Blotch in Almonds**

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

ALMOND		
Disease Suppression	Application Rate	Application Instructions
Red Leaf Blotch (RLB) ( <i>Polystigma amygdalinum</i> )	6.25 to 6.84 fl oz/acre  (0.203 – 0.222 lb/acre fluopyram)	<b>Soil Applications</b> - Apply specified dosage by chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment. For optimum results, apply to newly planted trees or those previously trained to drip, trickle or micro-sprinkler irrigation. Soil must be lightly pre-wetted to break soil surface tension prior to applications.  Minimum 30-day interval between soil applications.
<b>Restrictions:</b> <ul style="list-style-type: none"><li>Maximum single application rate: 6.84 fl oz/acre of VELUM ONE (0.222 lb/acre fluopyram).</li><li>Maximum annual application rate: 13.7 fl oz of VELUM ONE per acre (0.445 lb/acre fluopyram) per year.</li><li>Maximum number of applications per year: 2 (at 6.84 fl oz/acre of VELUM ONE).</li><li>Minimum soil retreatment interval: 30 days.</li><li><b>DO NOT</b> apply more than 0.446 lbs fluopyram per acre per year, regardless of formulation or method of application (soil or foliar).</li><li>Apply using chemigation equipment.</li><li><b>DO NOT</b> apply VELUM ONE within 14 days of harvest.</li><li>To limit the potential for development of disease resistance to this fungicide class, <b>DO NOT</b> make more than 2 sequential applications of VELUM ONE or any Group 7-containing fungicide before rotating with a fungicide from a different Group.</li><li>For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after VELUM ONE be a product from a different FRAC group.</li></ul>		

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

**Bayer CropScience LP**  
800 N. Lindbergh Blvd.  
St. Louis, MO 63167