

SUPPLEMENTAL LABELING

HALOSULFURON-METHYL	GROUP	2	HERBICIDE
---------------------	-------	---	-----------



Sandea® is a selective herbicide for control of listed weeds including nutsedge

This supplemental label expires on August 28, 2027, and must not be used or distributed after this date.

EPA REG. No. 81880-18-10163

ACTIVE INGREDIENT:	% BY WT.
Halosulfuron-methyl.....	75.0%
OTHER INGREDIENTS	25.0%
	TOTAL 100.0%

Contains 0.75 lb active ingredient per lb of product

KEEP OUT OF REACH OF CHILDREN CAUTION

Read the entire label before using this product.

- This labeling must be in the possession of the user at the time of pesticide application.
- It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

CROP	OZ/ACRE	DIRECTIONS FOR USE
GRASSES GROWN FOR SEED	3/4 - 1 1/3	<p>SPRING PLANTED TALL FESCUE GROWN FOR SEED WEST OF THE CASCADES</p> <p>For postemergence control of listed broadleaf weeds, apply 3/4 to 1 1/3 oz/A (0.035 to 0.062 lb ai/A). Apply as a broadcast spray in a minimum of 10 gallons of water/acre to new establishment seedling tall fescue in the spring once the first tiller of the crop is established.</p> <p>Applications for the control of sharppoint fluvellin must be made when the basal diameter of the weed is the size of a U.S. quarter, or smaller, and before stem elongation or runner formation.</p> <p>Tank mixing SANDEA with pyraflufen ethyl, pyrasulfotole, or saflufenacil and/or other herbicides will improve weed control.</p> <p>It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture.</p>
		<p>PRECAUTIONS:</p> <ul style="list-style-type: none"> • This product is effective if no rainfall occurs within 3 hours, but best results are obtained with no rainfall or irrigation for at least 8 hours. • Avoid application of SANDEA when grass seed crops or weeds are under stress since crop injury and poor weed control may result. • Refer to "Mixing Instructions", "Use Precautions", and "For Optimum Results" for detailed and important use information on the application of SANDEA. <p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply as an over the top spray to desirable shrubs or trees. • DO NOT exceed the Maximum Single Application Rate of 1 1/3 oz/A (0.062 lb ai/A). • DO NOT make more than 2 applications of SANDEA per year. • DO NOT apply more than 2 2/3 oz/A (0.125 lb ai/A) per year. • Minimum of 14 days between applications. • DO NOT apply by rope-wick wiper application.

EPA Reg. No. 81880-18-10163

NET CONTENTS _____

Distributed by:
Gowan Company, LLC
PO Box 5569
Yuma, AZ 85366-5569

Gowan®