

RESTRICTED USE PESTICIDE

May injure (phytotoxic) susceptible non-target plants. For retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. Commercial and certified applicators must ensure that all persons involved in these activities are informed of the precautionary statements.



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

CORVUS™ Herbicide

For Control of Broadleaf Weeds and Grasses in Corn

EPA Reg. No. 264-1066

SLN. No. SD-090002

24(c) Supplemental Label

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF SOUTH DAKOTA

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements and precautions on the EPA registered label. Do not apply this product through any type of irrigation system.

Environmental Precautions and Soil Type Restrictions

REFER TO THE CONTAINER LABEL FOR ADDITIONAL USE PRECAUTIONS AND DIRECTIONS

ENVIRONMENTAL PRECAUTIONS

AGRICULTURAL CHEMICALS HAVE THE POTENTIAL TO MOVE INTO SHALLOW GROUNDWATER. THE FOLLOWING RESTRICTIONS HAVE BEEN DEVELOPED TO PROTECT DRINKING WATER SUPPLIES.

In fields having soils with less than 15% field moisture holding capacity, special care must be taken not to over-irrigate, since substantial over-irrigation promotes the leaching of chemicals.

SOIL TYPE RESTRICTIONS

SOUTH DAKOTA

If the water table is less than 25 feet below ground surface, do not apply to the following vulnerable sandy loam, loamy sand or sand soils. If the water table is unknown, do not apply CORVUS™ Herbicide to a restricted soil.

Almeria	Dailey	Glendive	Minnewaukan	Talmo
Alwilda	Dix	Hamar	Munjor	Tassel
Anselmo	Doger	Hanly	O'Neill	Telfer
Arvilla	Dogiecreek	Hecla	Orton	Thurman
Assinniboine	Duda	Henkin variant	Orwet	Trembles
Aylmer	Dunday	Holt	Platte	Trey
Bankard	Dwyer	Hopdraw	Renshaw	Tryon
Banks	Eckley	Inavale	Reva	Tusler
Bantry	Egeland	Ipaga	Rhame	Tuthill
Beisigl	Ekalaka	Ladner	Riverwash	Twilight
Blackhall	Els	Las Animas	Rockoa	Ulen
Blendon	Elsmere	Lihen	Ronson	Valent
Boel	Evridge	Lute	Sarpy	Valentine
Butche	Fedora	Maddock	Schamber	Vebar
Cass	Flasher	Manning	Serden	Venlo
Chappell	Fleak	Manter	Seroco	Wabek
Chinook	Fossum	Marlake	Simeon	Whitelake
Claire	Gannett	Mawer	Sioux	Yecross
Cohagen	Glenberg	McCaffery	Stirum	Zeona
Craft	Glenberg variant	Meadin	Storla	

If a field contains several soil types, one of which is a vulnerable soil listed above, and the water table is less than 25 feet, do not apply CORVUS™ Herbicide. See reverse side for General Application Directions.

GENERAL APPLICATION DIRECTIONS

SPRAY DRIFT PRECAUTIONS

Controlling spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential of spray drift. The applicator and the grower are responsible for considering all these factors when making application decisions.

Do not apply this product using aerial application equipment.

Do not apply when wind conditions favor drift to non-target sites. To minimize spray drift to non-target areas:

Use low pressure application equipment capable of producing a large droplet spray. Do not use nozzles that produce fine droplet spray. Minimize drift by using sufficient spray volume to ensure adequate coverage with large droplet size sprays.

Keep ground driven spray boom as low as possible above the target surface.

Make application when the wind velocity favors on-target product deposition. Do not spray near sensitive plants if wind is gusty or in excess of 10 mph and moving in the direction of sensitive plants or sensitive areas.

Low humidity and high temperatures increase the likelihood of spray drift to sensitive areas. Avoid spraying during conditions of low humidity and/or high temperatures. Do not apply during inversion conditions.

MIXING, LOADING AND HANDLING PRECAUTIONS

Do not clean, wash, load, rinse, or empty equipment containing CORVUS™ Herbicide within 150 feet of any well, lake, stream, stream bed, or wetland unless it is done so over a South Dakota Department of Agriculture approved operational area containment.

Any spilled material must be immediately recovered. Discharges in excess of 25 pounds active ingredient must be reported to the department. Recovered material must be properly stored and handled until proper disposal or use has occurred. The owner or operator shall provide written documentation to the department describing the cause of the discharge, recovery and sampling procedures, analysis reports, and disposition of the recovered materials within 30 days following the date the discharge occurred.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at www.bayercropscienceus.com

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).