

Bayer CropScience LP 800 N. Lindbergh Blvd. St. Louis, Missouri 63167 1-866-99BAYER (1-866-992-2937)

## Luna Sensation®

Reduced application rates in grape EPA Reg. No. 264-1090

# FIFRA 2(ee) Recommendation

| FLUOPYRAM       | GROUP | 7  | FUNGICIDE |
|-----------------|-------|----|-----------|
| TRIFLOXYSTROBIN | GROUP | 11 | FUNGICIDE |

## FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF CALIFORNIA

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

#### **DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its label.

### Fruit, small vine climbing, except fuzzy kiwifruit, subgroup 13-07F

Amur River Grape; Gooseberry; Grape; Kiwifruit, Hardy; Maypop; Schisandra Berry; cultivars varieties, and/or hybrids of these.

**Note**: Do not apply or allow drift to **Concord grapes** or crop injury may occur. Rinse spray equipment before application of other products to **Concord grapes** or crop injury may occur.

| Disease Control   | Application Rate      | Application Instructions  |
|---|-----------------------|---|
| Powdery mildew (Uncinula necator)                       | 5.0 to 7.0 fl oz/acre | Apply using ground, aerial, or chemigation equipment.  Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14- to 21-day interval. When disease pressure is severe, use the shorter intervals. |
| Botrytis bunch rot /<br>Gray mold<br>(Botrytis cinerea) | 6.5 to 7.0 fl oz/acre | Apply using ground, aerial, or chemigation equipment.  Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 12- to 21-day interval. When disease pressure is severe, use the shorter intervals. |

| Fruit, small vine climbing, except fuzzy kiwifruit, subgroup 13-07F (continued)         |                          |   |  |
|---|--------------------------|---|--|
| Disease Control   | Application Rate         | Application Instructions  |  |
| Phomopsis cane and leaf spot (Phomopsis viticola)  Black rot (Guignardia bidwellii)     | 5.0 to 7.0<br>fl oz/acre | Apply using ground, aerial, or chemigation equipment.  Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14- to 21-day interval. When disease pressure is severe, use the shorter intervals. |  |
| Disease Suppression   | Application Rate         | Application Instructions  |  |
| Downy mildew ( <i>Plasmopara viticola</i> )  Aspergillus rot ( <i>Aspergillus</i> spp.) | 5.0 to 7.0 fl oz/acre    | Apply using ground, aerial, or chemigation equipment.  Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14- to 21-day interval. When disease pressure is severe, use the shorter intervals. |  |

#### **Restrictions:**

- Do not apply more than 21 fl oz of LUNA SENSATION (0.345 lb fluopyram and 0.345 lb trifloxystrobin) per acre per year.
- Do not apply more than 0.446 lbs fluopyram or 0.75 lbs trifloxystrobin per acre per year, including soil and foliar uses, regardless of formulation or method of application.
- Do not make more than 3 applications per year at 7.0 fl oz/acre of LUNA SENSATION, or 4 applications per year at 5.0 fl oz/acre of LUNA SENSATION.
- Do not apply LUNA SENSATION within 45 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, do not make more than 2 sequential applications of LUNA SENSATION or any Group 7 or Group 11 containing fungicide before rotating with a fungicide from a different Group.

For <u>MEDICAL</u> And <u>TRANSPORTATION</u> Emergencies <u>ONLY</u> Call 24 Hours A Day 1-800-334-7577. For <u>PRODUCT USE</u> Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions. For additional information call toll-free 1-866-99BAYER (1-866-992-2937).