



FLUOPYRAM	GROUP	7	FUNGICIDE
TEBUCONAZOLE	GROUP	3	FUNGICIDE

## FIFRA 2(ee) Recommendation

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF ARIZONA AND CALIFORNIA**

### LUNA EXPERIENCE®

EPA Reg. No. 264-1091

#### For Control of Red Leaf Blotch in Almonds

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

ALMOND		
Disease Control	Application Rate	Product Instructions
Red Leaf Blotch (RLB) ( <i>Polystigma amygdalinum</i> )	6.0 to 8.0 fl oz/acre  (0.078 – 0.104 lb/acre fluopyram) (0.078 – 0.104 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. For best results, make foliar applications up to 2 weeks post petal fall. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
<b>Restrictions:</b> <ul style="list-style-type: none"><li>Maximum single application rate: 8 fl oz/acre of LUNA EXPERIENCE (0.104 lb/acre fluopyram and 0.104 lb/acre tebuconazole).</li><li>Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year.</li><li>Maximum number of applications per year: 5 (at 6.0 fl oz/acre of LUNA EXPERIENCE)</li><li>Minimum retreatment interval: 7 days</li><li>Apply using ground, aerial, or chemigation equipment.</li><li>Regardless of formulation or method of application, <b>DO NOT</b> apply more than 0.446 lbs Fluopyram or 0.900 lbs tebuconazole per acre per year, including soil and foliar uses.</li><li><b>DO NOT</b> apply LUNA EXPERIENCE within 35 days of harvest.</li><li><b>DO NOT</b> cut cover crops in treated areas for feed, or allow livestock to graze treated areas.</li><li>To limit the potential for development of disease resistance to these fungicide classes, <b>DO NOT</b> make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.</li></ul>		

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

Bayer CropScience LP  
800 N. Lindbergh Blvd.  
St. Louis, MO 63167