

RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms. For retail sale to and use only by Certified Applicators, or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.



EPA REG. NO. 279-3313

2(ee) Recommendation

**FOR DISTRIBUTION AND USE IN
ALL REGISTERED STATES (Except Texas)**

**THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL
DIRECTIONS FOR USE WHICH ARE RECOMMENDED BY
FMC CORPORATION UNDER FIFRA SECTION 2(ee) AND HAS
NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.**

This recommendation for BRIGADE® 2EC Insecticide/Miticide is valid until 12/31/2030, or until withdrawn, canceled or suspended.

Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS
ON THE EPA REGISTERED BRIGADE® 2EC INSECTICIDE/MITICIDE (EPA REG. NO. 279-3313) LABEL MUST BE FOLLOWED.
THIS RECOMMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION.**

For control of spotted lanternfly in the listed crops.

CROP	RATE OF APPLICATION	RESTRICTIONS
Grape	6.4 fl. oz./A (0.10 lb ai/A)	<ul style="list-style-type: none">• DO NOT apply more than 6.4 fl. oz/A (0.10 lb ai/A) of product per application.• DO NOT apply more than 6.4 fl. oz/A (0.10 lb ai/A) of product per year.• DO NOT make more than 1 application per year.• DO NOT apply within 30 days of harvest.

Use Directions:

Thorough coverage is essential to achieve control.

Application by ground: Apply in a minimum of 25 gallons of finished spray per acre with ground equipment.

Application by air: Apply in a minimum of 10 gallons of finished spray per acre. When applying by air, 1 to 2 quarts of emulsified oil may be substituted for 1 to 2 quarts of water in the finished spray.

CROP	RATE OF APPLICATION	RESTRICTIONS
Hops*	6.4 fl. oz./A (0.10 lb ai/A)	<ul style="list-style-type: none">• DO NOT apply more than 6.4 fl. oz/A (0.10 lb ai/A) of product per application.• DO NOT apply more than 19.2 fl. oz/A (0.30 lb ai/A) of product per year.• Use of ultra low volume (ULV) application on hops is prohibited.• DO NOT make more than 3 applications per year.• DO NOT make applications less than 21 days apart.• DO NOT apply within 14 days of harvest.

Use Directions:

Application by ground: For best results, full coverage is essential. Early season recommendation is 100-150 gallons of finished spray per acre. Late season recommendation is 200-250 gallons of finished spray per acre.

*Not for use in CA unless accompanied by supplemental labeling.



FMC Corporation
2929 Walnut Street
Philadelphia, PA, 19103

CROP	RATE OF APPLICATION	RESTRICTIONS
Pears	6.4 – 12.8 fl. oz./A (0.10 – 0.20 lb ai/A)	<ul style="list-style-type: none"> • DO NOT apply more than 12.8 fl. oz/A (0.20 lb ai/A) of product per application. • DO NOT apply more than 32 fl. oz/A (0.50 lb ai/A) of product per year with no more than 28.8 fl. oz/A (0.45 lb ai/A) of product applied after petal fall. • DO NOT make more than 3 applications per year. • DO NOT graze livestock in treated orchards or cut treated cover crops for feed. • DO NOT make applications less than 30 days apart. • DO NOT apply within 14 days of harvest.
Use Directions: Application by ground: Apply as a dilute (minimum of 200 gallons of finished spray per acre) or concentrate (minimum of 50 gallons of finished spray per acre) spray in sufficient water to provide thorough coverage. Application by air: Apply the specified dosage in a minimum of 10 gallons of finished spray per acre by air.		

GENERAL DIRECTIONS FOR USE (ALL LISTED CROPS):

Thorough coverage is essential to achieve control.

Multiple applications may be required to maintain control and prevent reinfestation from migrating adults into the crop. Spotted lanternfly are mobile and may re-infest crops quickly. If another treatment is needed before the minimum retreatment interval, use another insecticide to maintain control.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or if bees are visiting the treatment area. **Protect pollinating insects by following label directions intended to minimize drift and to reduce risk to these organisms.**

The use of bifenthrin is prohibited in areas that may result in exposure of endangered species to bifenthrin. Prior to use in a particular county contact the local extension service for procedures and precautions to use to protect endangered species.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED BRIGADE® 2EC INSECTICIDE/MITICIDE (EPA REG. NO. 279-3313) LABEL MUST BE FOLLOWED. THIS RECOMMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION.

Brigade and FMC are trademarks of FMC Corporation or an affiliate.
©2025 FMC Corporation



FMC Corporation
2929 Walnut Street
Philadelphia, PA, 19103