



Supplemental Label

BIFENTURE® EC Agricultural Insecticide

(EPA Reg. No. 70506-57)

DIRECTIONS FOR USE ON LEAFY BRASSICAS (Crop Subgroup 5-B)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Follow all applicable directions, restrictions and precautions on the federally-registered label.

This label must be in the possession of the user at the time of pesticide application.

CROP	PESTS CONTROLLED	RATE BIFENTURE EC/ACRE	APPLICATION INSTRUCTIONS	PHI (days)
Broccoli raab Chinese cabbage (bok choy) Collards Kale Mizuna Mustard Greens Mustard Spinach Rape Greens	Aphids Armyworms Corn Earworm Crickets Cucumber Beetles Cutworms Diamondback Moth Flea Beetles Grasshoppers Ground Beetles Imported Cabbageworm Japanese Beetles (adults) Leafhoppers Loopers Saltmarsh Caterpillar Stink Bugs Thrips Tobacco Budworm Whitefly Wireworm (adults)	0.033-0.10 lb. ai (2.1-6.4 fl. oz. product)	Apply in water in a minimum of 10 gallons/A by ground or in a minimum of 2 gallons/A by air. For best control thorough coverage of foliage is necessary. When applied by air, 1 to 2 quarts of emulsified oil may be substituted for 1 to 2 quarts of water in the finished spray. Do not make applications less than 7 days apart. Do not apply more than 0.4 lb. a.i. /A per season.	7
	Banks Grass Mite Carmine Mite Lygus spp. Pacific Spider Mite Twospotted Spider Mite	0.08 - 0.10 lb. ai (5.12 - 6.4 fl. oz. product)		

Rev. 5/09
CA appr. 10/28/09

United Phosphorus, Inc.
630 Freedom Business Center
King of Prussia, PA 19406
1-800-438-6071 • www.upi-usa.com