

FIFRA Sec. 2(ee) Recommendation

Veltyma[®]

Fungicide

Reduced use rate for early season disease control in small grains grown in Idaho, Nevada, Oregon, Utah, and Washington

This Section 2(ee) Recommendation expires on December 31, 2027.

EPA Reg. No. 7969-409

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

All applicable directions, restrictions, precautions, and Conditions of Sale and Warranty on the EPA registered label are to be followed.

Information contained in this Section 2(ee) Recommendation is not intended to replace or amend any product labeling. Always read and follow all label directions when using any pesticide alone or in tank mix combinations.

The user must have this recommendation in their possession at the time of use.

Directions For Use

For early season control of tan spot, Septoria leaf and glume blotch, and spot blotch, apply 5 to 7 fl ozs per acre of **Veltyma[®] Fungicide** in combination with an herbicide application or when conditions favor disease development. When the 5 to 7 fl ozs early season application rate is used, a second application of fungicide may be required to protect the emerged flag leaf. Follow all use directions found on the **Veltyma** container label.

Veltyma is a registered trademark of BASF.

© 2024 BASF Agricultural Solutions US LLC
All rights reserved.

007969-00409.20240221.NVA 2024-04-0554-0035