## FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

FENPYROXIMATE | GROUP | 21A | INSECTICIDE



EPA Reg. No. 71711-57

Crop: Grape

Pest: Powdery mildew (Erysiphe necator)

States: California, Washington

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This labeling and the EPA approved container label must be in the possession of the user at the time of application. Read the label affixed to the container for FujiMite® 20SC Miticide/Insecticide before applying. Use of FujiMite 20SC Miticide/Insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for FujiMite 20SC Miticide/Insecticide.

## For Management of Powdery Mildew (Erysiphe necator) in Grape

- FujiMite 20SC Miticide/Insecticide is an effective contact powdery mildew fungicide. Direct contact with the fungal infection is necessary to achieve management of powdery mildew.
- FujiMite 20SC Miticide/Insecticide can be used to dessicate sporulating infections of powdery mildew on foliage and grape clusters.
- Apply FujiMite 20SC Miticide/Insecticide at 7.45 fl oz (0.105 lb ai) per acre.
- Make a foliar application using 200 to 300 gallons of water per acre or an adequate spray volume to obtain complete spray coverage of the infected foliage and fruit.
- It is recommended FujiMite 20SC Miticide/Insecticide be used with a known safe surfactant to improve spray coverage.
- To extend residual control, tank mix FujiMite 20SC Miticide/Insecticide with an effective systemic or translaminar powdery mildew fungicide.

©2025 Nichino America, Inc. FujiMite is a registered trademark of Nichino America, Inc.



Nichino America, Inc. 4550 Linden Hill Road, Suite 501 Wilmington, DE 19808 888-740-7700