

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## RESTRICTED USE PESTICIDE

May Injure (Phytotoxic) Susceptible, Non-Target Plants. For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. Commercial certified applicators must also ensure that all persons involved in these activities are informed of the precautionary statements.

## Tordon<sup>®</sup> 22K

EPA Reg. No. 62719-006

**For Control of Certain Troublesome Woody Plants on Rangelands, Permanent  
Grass Pastures and Non-Cropland Areas**

**REGISTERED USES OF TORDON 22K FOR THE STATE OF HAWAII  
(SALE AND USE RESTRICTED TO PICLORAM ANNUAL USE PERMIT HOLDERS)**

### NOTICE:

- **This supplemental labeling must be in the possession of the user at the time of application.**
- Before using Tordon 22K, read and carefully follow all applicable Use Directions, Use Precautions, Precautionary Statements and Storage and Disposal directions on the label affixed to the product container. Also, Read and carefully follow all Use Precautions in this Supplemental Labeling
- Before buying or using this product, read "**Warranty Limitations and Disclaimer**" on the Tordon 22K label. If terms are not acceptable, return unopened package at once to seller for refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under "**Warranty Limitations and Disclaimer**".
- Do not apply Tordon 22K through any type of irrigation system.
- Before applying, be certain that use of Tordon 22K conforms to all local regulations.

### Directions For Use

Use Tordon<sup>®</sup> 22K herbicide to control troublesome woody plants such as lantana, melastoma, guava, Java plum, pamakani, cat's claw, gorse, firebush and hau as well as many other woody plants, broadleaf weeds and vines infesting rangelands and permanent grass pastures and non-cropland area in the state of **Hawaii**. Treat anytime during the year when the plants are actively growing. Do not apply as a broadcast spray.

**Individual Plant Foliar Treatment:** Mix Tordon 22K at the rate of 2 quarts per 100 gallons of water. Add 2 quarts of surfactant to the spray mixture. Apply as a low pressure (10-30 psi) spray to thoroughly wet target vegetation, including leaves, stems and trunks of woody plants. For preparing smaller amounts of spray mixture, mix 2.5 fluid ounces of Tordon 22K and 2.5 fluid ounces of surfactant per 4 gallons of water and apply as above.

Re-treat in subsequent years as needed.

**Cut Surface Treatments:** For control of firebush, hau, Java plum and guava with trunks of 3 inches in diameter or larger. Use Tordon 22K diluted 1 to 4 in water as directed below.

**Tree Injector Application:** Application should be made by injecting 1 milliliter of the diluted herbicide solution through the bark at intervals of 3 inches between edges of the injection wound. The injections should completely surround the tree trunk at any convenient height.

**Frill or Girdle Application:** Make a single girdle through the bark completely around the tree trunk at a convenient height. Wet the cut surface with the diluted herbicide solution.

#### USE PRECAUTIONS

- **Do not graze dairy animals on treated areas** within two weeks after application. **Do not slaughter meat animals** grazing on treated areas until two weeks have elapsed after application.
- **Do not allow spray drift.** Tordon herbicides are highly active. Very small amounts may cause damage to plants if applied during either growth or dormant periods. Do not apply or otherwise permit Tordon 22K or sprays containing it to contact desirable plants such as pineapple, papaya, macadamia, coffee, vegetables, flowers, grapes, fruits trees, ornamentals, tomatoes, potatoes, beans of all types and other valuable broadleaf plants, nor soil containing roots of nearby valuable plants. Apply Tordon 22K only when there is little or no wind or no hazard from spray drift.
- **Do not contaminate water.** To avoid injury to crops or other desirable plants, do not contaminate irrigation ditches or water intended for irrigation or domestic purposes. Do not treat or allow spray drift to fall onto inner banks or bottom of irrigation ditches, either dry or containing water, or other channels that carry water that may be used for irrigation purposes.
- **Do not transfer livestock directly from treated areas onto broadleaf crop areas** without allowing 7 days on untreated grass pastures. Urine may contain enough picloram to cause crop injury.
- **Do not use manure from animals grazing treated areas** on land used for growing broadleaf crops, ornamentals, orchards or other valuable plants. Manure may contain enough picloram to cause crop injury.
- **Do not mix with other Pesticides** unless the pesticide is EPA approved for use on sites for which Tordon 22K use is approved. Use pesticide mixtures only in accordance with University of Hawaii treatment recommendations.
- **Cleaning Spray Equipment.** Carefully follow equipment cleaning instructions on Tordon 22K label.
- **Container Disposal.** Carefully follow container disposal instructions on Tordon 22K label.

®Trademark of Dow AgroSciences LLC

R111-004

Approved: 01/14/91

Replaces 122-42-001N