

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

FENPYROXIMATE GROUP 21A INSECTICIDE



EPA Reg. No. 71711-57

Crop: Citrus Fruit Group (Crop Group 10-10)

Australian desert lime; Australian finger-lime; Australian round lime; Brown River finger lime; calamondin; citron; citrus hybrids; grapefruit; Japanese summer grapefruit; kumquat; lemon; lime; Mediterranean mandarin; Mount White lime; New Guinea wild lime; orange, sour; orange, sweet; pummelo; Russell River lime; satsuma mandarin; sweet lime; Tachibana orange; Tahiti lime; tangelo; tangerine (mandarin); tangor; trifoliate orange; uniq fruit; cultivars, varieties, and/or hybrids of these

Pest: Citrus mealybug (*Planococcus citri*)

States: California, Arizona

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This labeling and the EPA approved container label must be in the possession of the user at the time of application. Read the label affixed to the container for FujiMite® 20SC Miticide/Insecticide before applying. Use of FujiMite 20SC Miticide/Insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for FujiMite 20SC Miticide/Insecticide.

For Control of Citrus Mealybug (*Planococcus citri*) in Citrus Fruits

- Apply FujiMite 20SC Miticide/Insecticide at 7.45 to 11.2 fl oz (0.105 to 0.158 lb ai) per acre.
- Apply by ground using 200 to 750 gallons of water per acre or an adequate spray volume to obtain complete spray coverage of the infested foliage and fruit.
- Maintain a minimum Fujimite 20SC Miticide/Insecticide concentration of at least 3.73 fl oz per 250 GPA.
 - If using 500 GPA, a minimum volume of Fujimite 20SC Miticide/Insecticide at 7.45 fl oz/A is recommended.
 - If using 750 GPA, a minimum volume of Fujimite 20SC Miticide/Insecticide at 11.2 fl oz/A is recommended.
- It is recommended FujiMite 20SC Miticide/Insecticide be used with a safe surfactant or 415 oil to improve spray coverage.
- FujiMite 20SC Miticide/Insecticide is effective on all mealybug stages including adults; however, application should be made when most of the population is at the first or second instars.
- FujiMite 20SC Miticide/Insecticide is a contact insecticide on mealybugs; therefore, direct contact with the insect is necessary to achieve maximum control.
- For best results, tractor speed should not exceed 1.5 mph.

©2025 Nichino America, Inc. FujiMite is a registered trademark of Nichino America, Inc.



Nichino America, Inc.
4550 Linden Hill Road, Suite 501
Wilmington, DE 19808
888-740-7700