

BIOFUNGICIDE

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U.S. Environmental Protection Agency.

Double Nickel 55TM Biofungicide

Active Ingredient:

Bacillus amyloliquefaciens strain D747*	25.0%
Other Ingredients:	75.0%
Total	. 100.0%
*Contains a minimum of 5×10 ¹⁰ colony-forming units (cfu) per gram	

EPA Reg. No. 70051-108

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PEST or DISEASE:

For suppression of *Phytophthora* and *Fusarium* root rots in grapes (including wine grapes, table grapes, and raisins) and almonds.

APPLICATION INSTRUCTIONS:

Begin applications in early spring, timed for root flush and early shoot growth. Continue applications on a 4-6 week reapplication interval with a final application timed for fall root flush. Apply via drip (elevated, buried or ground-lay), banded drench, microsprinkler, or other common chemigation techniques.

All applicable directions, application rates, restrictions, and precautions on the EPA-registered labels are to be followed. This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the cautionary statements plus all other information appearing on the product labels. This recommendation must be in the possession of the user at the time of pesticide application.