



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

SIVANTO™ Prime

Reduce rate for foliar applications in Legume Vegetables for control of Soybean Aphid.

EPA Reg. No. 264-1141

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF MINNESOTA, NORTH DAKOTA, SOUTH DAKOTA, AND WISCONSIN

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

LEGUME VEGETABLES (SUCCULENT OR DRIED) – FOLIAR

Crops of Crop Group 6, including: Edible Podded and Succulent Shelled Pea and Bean and Dried Shelled Pea and Bean Bean (*Lupinus* spp., including grain lupin, sweet lupin, white lupin, and white sweet lupin), Bean (*Phaseolus* spp., including field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, and wax bean), Bean (*Vigna* spp., including adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, Southern pea, urd bean, and yardlong bean), Pea (*Pisum* spp. including dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, and sugar snap pea), Other Beans and Peas (Broad bean (fava bean), Chickpea (garbanzo bean), Guar, Jackbean, Lablab bean (hyacinth bean), Lentil, Pigeon pea, soybean (immature seed), and Sword bean), Soybean

Pests Controlled	Product Rate (fl oz/A)
Soybean aphid	5.0

Foliar Application Restrictions:

Pre-Harvest Interval (PHI): **7 days** – Forage, Leaves, Vines, Pods, Cutting for Hay, or Seed (fresh or dry, except dry soybean seed); **21 days** – dry soybean seed

Minimum interval between applications: **10 days**

Minimum application volumes: **10 gallons/Acre** (Ground); **3 gallons/Acre** (Aerial)

Do not apply more than **0.365 lb** flupyradifurone (**28.0 fl oz** of SIVANTO prime) per acre per calendar year on legume vegetables (succulent or dried), regardless of product or formulation.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*